

HEAT KERNEL ESTIMATES FOR THE GRUŠIN OPERATOR

MARTIN PAULAT

ABSTRACT. We study the geometry associated to the Grušin operator

$$G = \Delta_x + |x|^2 \partial_u^2 \text{ on } \mathbb{R}_x^n \times \mathbb{R}_u,$$

to obtain heat kernel estimates for this operator. The main work is to find the shortest geodesics connecting two given points in $\mathbb{R}^{n+1}$. This gives the Carnot-Carathéodory distance d_{CC} , associated to this operator. The main result in the second part is to give Gaussian bounds for the heat kernel K_t in terms of the Carnot-Carathéodory distance. In particular we obtain the following estimate

$$|k_t(\zeta, \eta)| \leq C t^{-\frac{n}{2}-1} \min \left(1 + \frac{d_{CC}(\zeta, \eta)}{|x + \xi|}, 1 + \frac{d_{CC}(\zeta, \eta)^2}{4t} \right)^\alpha e^{-\frac{1}{4t} d_{CC}(\zeta, \eta)^2}$$

for all $\zeta = (x, u_1)$, $\eta = (\xi, u) \in \mathbb{R}^{n+1}$, where $\alpha = \max \left\{ \frac{n}{2} - 1, 0 \right\}$. Here the homogeneous dimension is $q = n + 2$, so that $\frac{n}{2} - 1 = \frac{q-4}{2}$. This shows that our result for $n \geq 2$ corresponds with the result on the Heisenberg group, which was given by Beals, Gaveau, Greiner in [1].

Acknowledgement. I would like to express my gratitude to my advisor Professor Dr. Detlef Müller for constant support and numberless helpful suggestions.

1. INTRODUCTION

The purpose of this article is to study the geometric properties of the Grušin operator

$$G = \Delta_x + |x|^2 \partial_u^2 \text{ on } \mathbb{R}_x^n \times \mathbb{R}_u$$

to give estimates for the heat kernel of this operator. One may write G as

$$G = \sum_{j=1}^n (X_j^2 + U_j^2),$$

with the smooth vector fields

$$X_j := \partial_{x_j}, \quad U_j := x_j \partial_u, \quad j = 1, \dots, n.$$

Note that G is hypoelliptic, since G satisfies the Hörmander condition. We are interested in the Carnot-Carathéodory distance associated to these vector fields, i.e. the length of a minimizing horizontal curve. A horizontal curve is a curve, whose tangents are a linear combination of the vector fields X_j , U_j , $j = 1, \dots, n$. Depending on the start and end point of the curve, there are a different number of locally minimizing curves, i.e. geodesics. In addition to the results of Calin, Chang, Greiner and Kannai in [3], we can compare their lengths to give explicit formulas for the Carnot-Carathéodory distance. The geodesics starting in $0 \in \mathbb{R}^{n+1}$ are very similar to those on the Heisenberg group (see [1]). But comparing the lengths of geodesics starting in a generic point $(x, 0) \in \mathbb{R}^n \times \mathbb{R}$ gets more complicated. Another difference occurs when studying curves connecting $(x, 0)$ and $(\pm x, u)$, where one gets two different types of geodesics (see Theorem (2.4) and (2.5)).

These results on the Carnot-Carathéodory distance allow us to give some point-wise estimates of the heat kernels.

Due to the fact that the partial Fourier transform of G in u leads to the rescaled Hermite operators, it is easy to calculate the heat kernel using Mehler's formula, up to the partial Fourier transform. One may also observe that G is translation invariant in the u -variable. This allows us to write the heat kernel in the form $K_t(x, \xi, u)$, $x, \xi \in \mathbb{R}^n$, $u \in \mathbb{R}$. So the solution to the heat equation

$$\begin{cases} (\partial_t - G)u = 0 \\ u(0, \cdot) = f \end{cases}$$

is given by

$$u(t, x) = \int_{\mathbb{R}^n} \int_{\mathbb{R}} K_t(x, \xi, u - \lambda) f(\xi, \lambda) d\lambda d\xi.$$

We are able to give global estimates for the kernel of the form

$$|K_t(x, \xi, u)| \leq F(x, \xi, u, t) e^{-\frac{1}{4t} d_{CC}((x, 0), (\xi, u))^2},$$

where F is a function with polynomial growth. It turns out that

$$F(x, \xi, u, t) \leq C t^{-\frac{n}{2}-1} \min \left(1 + \frac{d_{CC}(\zeta, \eta)}{|x + \xi|}, 1 + \frac{d_{CC}(\zeta, \eta)^2}{4t} \right)^\alpha,$$

with $\alpha = \max \{ \frac{n}{2} - 1, 0 \}$ and $C > 0$ is a constant independent of x, ξ, u, t . In particular our main result is the following

Theorem. For $\zeta = (x, 0)$, $\eta = (\xi, u) \in \mathbb{R}^{n+1}$ we have

$$(1.1) \quad |K_t(x, \xi, u)| \lesssim t^{-\frac{n}{2}-1} \min \left(1 + \frac{d_{CC}(\zeta, \eta)}{|x + \xi|}, 1 + \frac{d_{CC}(\zeta, \eta)^2}{4t} \right)^\alpha e^{-\frac{1}{4t} d_{CC}(\zeta, \eta)^2},$$

with $\alpha = \max \{ \frac{n}{2} - 1, 0 \}$.

For a better understanding of the exponent α , consider the dilation

$$\delta_r(x, u) := (rx, r^2u), \quad x \in \mathbb{R}^n, u \in \mathbb{R},$$

for $r > 0$. Then G is homogeneous with degree 2, and also the Carnot-Carathéodory distance is homogeneous with degree 2, i.e.

$$d_{CC}(\delta_r(\zeta), \delta_r(\eta)) = r^2 d_{CC}(\zeta, \eta), \quad \zeta, \eta \in \mathbb{R}^{n+1}.$$

In this setting the homogeneous dimension is $q = n + 2$, and $\frac{n}{2} - 1 = \frac{q-4}{2}$, $\frac{n}{2} + 1 = \frac{q}{2}$. This shows now that our result exactly coincides, in the case $n \geq 2$, with those on the Heisenberg group obtained by Beals, Gaveau, Greiner in [1].

There are some Gaussian-type estimates for heat kernels with greater generality. Sikora, for example, shows in [8]

$$|K_t(x, \xi, u)| \lesssim t^{-\frac{q}{2}} \left(1 + \frac{d_{CC}(\zeta, \eta)}{4t} \right)^{\frac{q-1}{2}} e^{-\frac{d_{CC}(\zeta, \eta)^2}{4t}}.$$

The methods that are used are completely different. The explicit formulas in our situation allow us to give a smaller exponent of the polynomial factor, i.e. $\frac{q-4}{2}$ instead of $\frac{q-1}{2}$.

2. THE SUB-RIEMANNIAN GEOMETRY ASSOCIATED TO THE GRUŠIN OPERATOR

Let M be a connected C^∞ -Manifold with smooth real vector fields $X_1, \dots, X_m$. For $x \in M$ and $v \in T_x M$ define:

$$(2.1) \quad \|v\|_x^2 := \inf \left\{ \sum_{j=1}^m \alpha_j^2 : v = \sum_{j=1}^m \alpha_j X_j(x) \right\},$$

where we use the convention, that $\inf \emptyset := \infty$, i.e. if v is not in the linear span of $X_1, \dots, X_m$, then we set $\|v\|_x^2 := \infty$. A horizontal curve $\gamma : [0, 1] \rightarrow M$ is an absolutely continuous curve such that $\dot{\gamma}(t) \in \text{span}\{X_1(\gamma(t)), \dots, X_m(\gamma(t))\}$ for almost every $t \in [0, 1]$. We define the length of a horizontal curve by

$$(2.2) \quad L(\gamma) := \int_0^1 \|\dot{\gamma}(t)\|_{\gamma(t)} dt.$$

Then the Carnot-Carathéodory distance of two points $p, q \in M$ associated to $X_1, \dots, X_m$ is defined by

$$(2.3) \quad d_{CC}(p, q) := \inf L(\gamma),$$

where the infimum is taken over all horizontal curves γ connecting p and q , i.e. $\gamma(0) = p$, $\gamma(1) = q$.

It is well known by Chow's theorem (see [4]) that, if $X_1, \dots, X_m$ and their brackets span the tangent space $T_x M$ at every point x of M , then any two points can be joined by such a curve, so that $d_{CC}(p, q) < \infty$, for any $p, q \in M$.

Now let $M = \mathbb{R}^{n+1}$ and

$$X_j = \partial_{x_j}, \quad U_j = x_j \partial_u, \quad j = 1, \dots, n.$$

Then the Grušin operator $G = \Delta_x + |x|^2 \partial_u^2$ reads in terms of these vector fields

$$G = \sum_{j=1}^n (X_j^2 + U_j^2).$$

The vector fields $X_1, \dots, X_n, U_1, \dots, U_n$ span the tangent space everywhere except along the line $|x| = 0$. But, since $[X_j, U_j] = \partial_u$, $j = 1, \dots, n$, all conditions are fulfilled to connect any two points by an absolutely continuous curve with finite length. If $\gamma : [0, 1] \rightarrow \mathbb{R}^{n+1}$ is an absolutely continuous curve, one has

$$\begin{aligned} \dot{\gamma} &= \dot{\gamma}_{n+1} \partial_u + \sum_{j=1}^n \dot{\gamma}_j \partial_{x_j} \\ &= \dot{\gamma}_{n+1} \partial_u + \sum_{j=1}^n \dot{\gamma}_j X_j \end{aligned}$$

almost everywhere. Since the vector fields U_j , $j = 1, \dots, n$ are not linear independent, we may write $\dot{\gamma}_{n+1} \partial_u$ as a linear combination of $U_1, \dots, U_n$ in different ways. So

$$\|\dot{\gamma}(t)\|_{\gamma(t)}^2 = \sum_{j=1}^n \dot{\gamma}_j(t)^2 + \inf \left\{ \sum_{j=1}^n \alpha_j^2 : \dot{\gamma}_{n+1}(t) = \sum_{j=1}^n \alpha_j \dot{\gamma}_j(t) \right\}.$$

To minimize this convex functional, one can use the method of Langrange multiplier, which gives

$$\|\dot{\gamma}(t)\|_{\gamma(t)}^2 = \frac{\dot{\gamma}_{n+1}(t)^2}{\sum_{j=1}^n \dot{\gamma}_j(t)^2} + \sum_{j=1}^n \dot{\gamma}_j(t)^2.$$

Now let $\gamma = (\gamma_{(1)}, \gamma_{(2)})$, with $\gamma_{(1)} = (\gamma_1, \dots, \gamma_n)$, $\gamma_{(2)} = \gamma_{n+1}$, then

$$\|\dot{\gamma}(t)\|_{\gamma(t)}^2 = \frac{\dot{\gamma}_{(2)}(t)^2}{|\gamma_{(1)}(t)|^2} + |\dot{\gamma}_{(1)}(t)|^2.$$

To calculate the Carnot-Carathéodory distance, one can minimize the “energy” integral

$$(2.4) \quad \int_0^1 \|\dot{\gamma}(t)\|_{\gamma(t)}^2 dt = \int_0^1 \left(\frac{\dot{\gamma}_{(2)}(t)^2}{|\gamma_{(1)}(t)|^2} + |\dot{\gamma}_{(1)}(t)|^2 \right) dt.$$

This leads to the Euler-Lagrange equations, which gives us local minimizing curves, i.e. geodesics:

$$(2.5a) \quad \ddot{\gamma}_{(1)} + \frac{\dot{\gamma}_{(2)}^2}{|\gamma_{(1)}|^4} \gamma_{(1)} = 0$$

$$(2.5b) \quad \frac{d}{dt} \frac{\dot{\gamma}_{(2)}}{|\gamma_{(1)}|^2} = 0.$$

Solving these equations yields the following result (see [6, 3, 7]).

All geodesics γ starting in $(x_1, u_1) \in \mathbb{R}^n \times \mathbb{R}$ are given by $\gamma^{b,c} = (\gamma_{(1)}^{b,c}, \gamma_{(2)}^{b,c}) : [0, 1] \rightarrow \mathbb{R}^n \times \mathbb{R}$, where

$$(2.6a) \quad \gamma_{(1)}^{b,c}(t) = \frac{c}{b} \sin(bt) + x_1 \cos(bt)$$

$$(2.6b) \quad \gamma_{(2)}^{b,c}(t) = \frac{|c|^2}{b} \left(\frac{t}{2} - \frac{\sin(2bt)}{4b} \right) + \frac{x_1 \cdot c}{b} \sin^2(bt) + |x_1|^2 b \left(\frac{t}{2} + \frac{\sin(2bt)}{4b} \right) + u_1$$

with parameters $b \in \mathbb{R}$, $b \neq 0$, $c \in \mathbb{R}^n$, and

$$(2.7) \quad \gamma^{0,c}(t) = (ct + x_1, u_1)$$

which is the limiting case $b \rightarrow 0$.

The length is given by

$$(2.8) \quad L(\gamma^{b,c}) = \sqrt{|c|^2 + |x_1|^2 b^2}.$$

To calculate the Carnot-Carathéodory distance, we are interested in the shortest geodesic joining two given points $(x_1, u_1), (x, u) \in \mathbb{R}^n \times \mathbb{R}$.

In our further analysis we will assume that $u_1 = 0$, since γ is a geodesic connecting $(x_1, 0)$, (x, u) , if and only if $\gamma + (0, u_1)$ is a geodesic connecting (x_1, u_1) , $(x, u + u_1)$; with the same length.

We will also assume that $u \geq 0$, since $\gamma_{(1)}^{b,c} = \gamma_{(1)}^{-b,c}$ and $\gamma_{(2)}^{b,c} = -\gamma_{(2)}^{-b,c}$.

Furthermore observe, that given $b \in \mathbb{R} \setminus \pi\mathbb{Z}^*$ the parameter $c \in \mathbb{R}^n$ is uniquely determined by $c = \frac{b}{\sin b} (x - x_1 \cos b)$, due to the boundary conditions. We will denote this geodesic by γ^b for short.

Given $b \in \pi\mathbb{Z}^*$ (note that this means $x = \pm x_1$), the parameter $c \in \mathbb{R}^n$ is determined by the whole sphere $|c| = \sqrt{2bu - |x_1|^2 b^2}$. But any $c \in \mathbb{R}^n$ with $|c| = \sqrt{2bu - |x_1|^2 b^2}$ will give the same length.

Theorem 2.1. *Given $(x_1, 0), (x, 0) \in \mathbb{R}^n \times \mathbb{R}$, the only geodesic connecting them is given by*

$$(2.9) \quad \gamma^0(t) = (x_1 + t(x - x_1), 0)$$

and its length, which is equal to the Carnot-Carathéodory distance, is

$$(2.10) \quad L(\gamma^0) = d_{CC}((x_1, 0), (x, 0)) = |x - x_1|.$$

Proof. We have to prove that $b = 0$ is the only parameter that allows connecting $(x_1, 0)$ and $(x, 0)$. But from the Euler-Lagrange equation, we know that $\dot{\gamma}_{(2)} = b |\gamma_{(1)}|^2$. That means $\dot{\gamma}_{(2)}(t) \geq 0$ or $\dot{\gamma}_{(2)}(t) \leq 0$ for all $t \geq 0$. But since we want $\gamma_{(2)}(0) = 0$ and $\gamma_{(2)}(1) = 0$, we conclude that $\dot{\gamma}_{(2)} = 0$, so that $b = 0$, hence $c = x - x_1$. $\square$

Theorem 2.2. *Given $u > 0$, there are infinitely many geodesics $\gamma^{m\pi, c_m}$, $m \in \mathbb{N}$, connecting $(0, 0) \in \mathbb{R}^n \times \mathbb{R}$ and $(0, u) \in \mathbb{R}^n \times \mathbb{R}$ given by (2.6) with $c_m \in \mathbb{R}^n$, $|c_m| = \sqrt{2m\pi u}$. Their lengths are*

$$(2.11) \quad L(\gamma^{m\pi, c_m}) = \sqrt{2m\pi u}.$$

In particular the shortest geodesics in this case, which give the Carnot-Carathéodory distance, are γ^{π, c_1} , $|c_1| = \sqrt{2\pi u}$; so

$$(2.12) \quad d_{CC}((0, 0), (0, u)) = \sqrt{2\pi u}.$$

Proof. The boundary conditions $\gamma_{(1)}^{b,c}(1) = 0$ and $\gamma_{(2)}^{b,c}(1) = u$ are equivalent to

$$0 = \frac{c}{b} \sin b \quad \text{and} \quad u = \frac{|c|^2}{2b} \left(1 - \frac{\sin(2b)}{2b}\right).$$

Since $u \neq 0$, we have $c \neq 0$. But then there exists $m \in \mathbb{N}$ with $b = m\pi$, so that $|c|^2 = 2m\pi u$. This gives the claim. $\square$




FIGURE 1. Geodesics starting at $(0, 0)$ and ending in $(0, u)$ with different parameters.

Now we consider the case where we want to connect $(x_1, 0)$ and (x, u) with $x \neq \pm x_1$, $u > 0$. The first lemma is just a rewritten formulation of the boundary conditions $\gamma^{b,c}(0) = (x_1, 0)$ and $\gamma^{b,c}(1) = (x, u)$.

Lemma 2.1. *Suppose $(x_1, 0), (x, u) \in \mathbb{R}^n \times \mathbb{R}$, $x \neq \pm x_1$, $u > 0$. Then $\gamma^{b,c}$ given by (2.6) is a geodesic connecting these two points, iff*

$$c = \frac{b}{\sin b} x - x_1 b \cot b$$

and b is any solution of

$$(2.13) \quad \frac{2u}{|x_1|^2 + |x|^2} = \frac{b}{\sin^2 b} - \cot b + \frac{2x \cdot x_1}{|x_1|^2 + |x|^2} \frac{1 - b \cot b}{\sin b}.$$

The square of the length of $\gamma^b = \gamma^{b,c}$ is then given by

$$(2.14a) \quad L^2(\gamma^{b,c}) = \frac{b^2}{\sin^2 b} \left(|x_1|^2 + |x|^2 - 2x \cdot x_1 \cos b \right)$$

$$(2.14b) \quad = 2bu + \left(|x_1|^2 + |x|^2 \right) b \cot b - 2x \cdot x_1 \frac{b}{\sin b}.$$

Proof. To find those geodesics connecting two points $(x_1, 0)$ and (x, u) , we have to find $b \neq 0$ and $c \in \mathbb{R}^n$, such that $\gamma_{(1)}^{b,c}(1) = x$ and $\gamma_{(2)}^{b,c}(1) = u$. Since $b \in \pi\mathbb{Z}$ means $x = \pm x_1$, we have that $b \notin \pi\mathbb{Z}$. Now $\gamma_{(2)}^{b,c}(1) = u$ gives:

$$(2.15) \quad \begin{aligned} 2u &= \frac{|c|^2}{b} \left(1 - \frac{\sin(2b)}{2b} \right) + 2 \frac{x_1 \cdot c}{b} \sin^2(b) + |x_1|^2 b \left(1 + \frac{\sin(2b)}{2b} \right) \\ &= \frac{1}{b} \left(|c|^2 + |x_1|^2 b^2 \right) - \frac{\sin(2b)}{2b^2} \left(|c|^2 - |x_1|^2 b^2 \right) + 2 \frac{x_1 \cdot c}{b} \sin^2 b \end{aligned}$$

And $\gamma_{(1)}^{b,c}(1) = x$ gives

$$(2.16) \quad \frac{b}{\sin b} x = c + x_1 b \cot b,$$

so that

$$(2.17) \quad \begin{aligned} \frac{b^2}{\sin^2 b} |x|^2 &= |c|^2 + 2c \cdot x_1 b \cot b + |x_1|^2 b^2 \cot^2 b \\ &= |c|^2 + 2c \cdot x_1 b \cot b - |x_1|^2 b^2 + \frac{|x_1|^2 b^2}{\sin^2 b} \end{aligned}$$

and

$$(2.18) \quad \begin{aligned} |c|^2 &= \frac{b^2}{\sin^2 b} |x|^2 - 2x_1 \cdot x \frac{b^2 \cot b}{\sin b} + |x_1|^2 b^2 \cot^2 b \\ &= \frac{b^2}{\sin^2 b} \left(|x|^2 + |x_1|^2 \right) - 2x_1 \cdot x \frac{b^2 \cot b}{\sin b} - |x_1|^2 b^2 \\ &= \frac{b^2}{\sin^2 b} \left(|x|^2 + |x_1|^2 - 2x \cdot x_1 \cos b \right) - |x_1|^2 b^2. \end{aligned}$$

We can continue with (2.15) by using equation (2.17)

$$\begin{aligned} 2u &= \frac{1}{b} \left(|c|^2 + |x_1|^2 b^2 \right) - \cot b \left(|x|^2 - |x_1|^2 \right) + 2 \frac{c \cdot x_1}{b} \cos^2 b + 2 \frac{c \cdot x_1}{b} \sin^2 b \\ &= \frac{1}{b} \left(|c|^2 + |x_1|^2 b^2 \right) - \cot b \left(|x|^2 - |x_1|^2 \right) + 2 \frac{c \cdot x_1}{b} \\ &= \frac{1}{b} \left(|c|^2 + |x_1|^2 b^2 \right) - \cot b \left(|x|^2 + |x_1|^2 \right) + 2 \frac{c \cdot x_1 + |x_1|^2 b \cot b}{b}. \end{aligned}$$

Now use equation (2.16) and (2.18)

$$\begin{aligned} 2u &= \frac{1}{b} \left(|c|^2 + |x_1|^2 b^2 \right) - \cot b \left(|x|^2 + |x_1|^2 \right) + 2 \frac{x \cdot x_1}{\sin b} \\ &= \left(\frac{b}{\sin^2 b} - \cot b \right) \left(|x|^2 + |x_1|^2 \right) + 2x \cdot x_1 \frac{1 - b \cot b}{\sin b}. \end{aligned}$$

And the square of the length is given by

$$\begin{aligned} L^2(\gamma^{b,c}) &= \frac{b^2}{\sin^2 b} \left(|x|^2 + |x_1|^2 - 2x \cdot x_1 \cos b \right) \\ &= 2bu + b \cot b \left(|x|^2 + |x_1|^2 \right) - 2 \frac{b}{\sin b} x \cdot x_1. \end{aligned}$$

□

Definition. For $b \in \mathbb{R} \setminus \pi\mathbb{Z}^*$, $-1 \leq a \leq 1$ define:

$$(2.19) \quad \mu(b) := \mu(b, a) := \frac{b}{\sin^2 b} - \cot b + a \frac{1 - b \cot b}{\sin b}$$

$$(2.20) \quad l(b) := l(b, a) := \frac{b^2}{\sin^2 b} (1 - a \cos b).$$

In the following we will often use the abbreviation

$$R = \sqrt{|x|^2 + |x_1|^2} \text{ and } a = \frac{2x_1 \cdot x}{|x_1|^2 + |x|^2}.$$

Next we will study the functions μ and $\tilde{\mu}$, $\hat{\mu}$, which will be introduced in the next lemma. (Note that $\hat{\mu}$ in this section does not mean Fourier transform!)

Lemma 2.2. For $b \in \mathbb{R} \setminus \pi\mathbb{Z}^*$, $0 \leq a \leq 1$, μ is a convex combination, i.e.

$$(2.21) \quad \mu(b) = (1 - a)\tilde{\mu} + a\hat{\mu}\left(\frac{b}{2}\right),$$

and for $-1 \leq a < 0$:

$$(2.22) \quad \mu(b) = (1 + a)\tilde{\mu}(b) - a\tilde{\mu}\left(\frac{b}{2}\right),$$

where

$$(2.23) \quad \tilde{\mu}(b) := \frac{b}{\sin^2 b} - \cot b, \quad b \in \mathbb{R} \setminus \pi\mathbb{Z}^*,$$

$$(2.24) \quad \hat{\mu}(b) := \frac{b}{\cos^2 b} + \tan b, \quad b \in \mathbb{R} \setminus \left\{ \left(k + \frac{1}{2}\right)\pi : k \in \mathbb{Z} \right\}.$$

Proof. For $b \in \mathbb{R} \setminus \pi\mathbb{Z}^*$:


$$\begin{aligned} \mu(b) &= (1 - a)\tilde{\mu}(b) + a \frac{1 - b \cot b + \frac{b}{\sin b} - \cos b}{\sin b} \\ &= (1 - a)\tilde{\mu}(b) + 2a \frac{\sin^2 \frac{b}{2} \left(1 + \frac{b}{\sin b}\right)}{\sin b} \\ &= (1 - a)\tilde{\mu}(b) + a \left(\frac{\frac{b}{2}}{\cos^2 \frac{b}{2}} + \tan \frac{b}{2} \right) \\ &= (1 - a)\tilde{\mu}(b) + a\hat{\mu}\left(\frac{b}{2}\right) \end{aligned}$$

and

$$\begin{aligned} \mu(b) &= (1 + a)\tilde{\mu}(b) - a \frac{b \cot b - 1 + \frac{b}{\sin b} - \cos b}{\sin b} \\ &= (1 + a)\tilde{\mu}(b) - 2a \frac{\cos^2 \frac{b}{2} \left(\frac{b}{\sin b} - 1\right)}{\sin b} \\ &= (1 + a)\tilde{\mu}(b) - a \left(\frac{\frac{b}{2}}{\sin^2 \frac{b}{2}} - \cot \frac{b}{2} \right) \\ &= (1 + a)\tilde{\mu}(b) - a\tilde{\mu}\left(\frac{b}{2}\right) \end{aligned}$$

□

Remark. Observe that all these functions $\mu, \tilde{\mu}, \hat{\mu}$ are odd. The following lemmata will show that $\tilde{\mu}|_{[0, \infty)} \geq 0$, $\hat{\mu}|_{[0, \infty)} \geq 0$ and therefore $\mu|_{[0, \infty)} \geq 0$. Since we assumed that $u \geq 0$, we only need information of these functions for $b \geq 0$.



The function $\tilde{\mu}$, which also appears in the study of geodesics on the Heisenberg group, where intensively studied by Beals, Gaveau, Greiner in [1]. We will use the following result:

Lemma 2.3. *The function $\tilde{\mu}$ is a monotone increasing diffeomorphism of the interval $[0, \pi)$ onto $[0, \infty)$. On each interval $(m\pi, (m+1)\pi)$, $m \in \mathbb{N}$, $\tilde{\mu}$ has a unique critical point $\tilde{b}_m$, which is a minimum. $\tilde{b}_m$ is implicitly given as the solution of the equation $1 - b \cot b = 0$ in the corresponding interval. On this interval, $\tilde{\mu}$ decreases strictly from $+\infty$ to $\tilde{\mu}(\tilde{b}_m)$ and then increases strictly to $+\infty$. Moreover,*

$$(2.25) \quad \tilde{\mu}(\tilde{b}_m) + \pi < \tilde{\mu}(\tilde{b}_{m+1}),$$

and

$$(2.26) \quad \tilde{\mu}(\tilde{b}_m) > m\pi.$$

The function $\hat{\mu}$ has been studied by Calin, Chang, Greiner, Kannai in [3]:

Lemma 2.4. *The function $\hat{\mu}$ is a monotone increasing diffeomorphism of $[0, \frac{\pi}{2})$ onto $[0, \infty)$. On each interval $(m\pi + \frac{\pi}{2}, (m+1)\pi + \frac{\pi}{2})$, $m \in \mathbb{N}_0$, $\hat{\mu}$ has a unique critical point $\hat{b}_m$, which is a minimum. $\hat{b}_m$ is implicitly given as the solution of the equation $1 + b \tan b = 0$ in the corresponding interval. On this interval, $\hat{\mu}$ decreases strictly from $+\infty$ to $\hat{\mu}(\hat{b}_m)$ and then increases strictly to $+\infty$. Moreover,*

$$(2.27) \quad \hat{\mu}(\hat{b}_m) \geq \pi \left(m + \frac{1}{2} \right).$$

Remark. Moreover $\tilde{\mu}''(b) > 0$, $b \notin \pi\mathbb{N}$ and $\hat{\mu}''(b) > 0$, $b - \frac{\pi}{2} \notin \mathbb{N}_0$, i.e. $\tilde{\mu}$ is strictly convex on each interval $(m\pi, (m+1)\pi)$, $m \in \mathbb{N}$ and $\hat{\mu}$ is strictly convex on each interval $(m\pi + \frac{\pi}{2}, (m+1)\pi + \frac{\pi}{2})$, $m \in \mathbb{N}_0$. This is another result of [1, 3]. Hence μ , as a convex combination, is also strictly convex on each interval $(m\pi, (m+1)\pi)$, $m \in \mathbb{N}$.

Combining the previous two lemmata we get the following result for the function μ :

Lemma 2.5. *Let $-1 < a < 1$. The function μ is monotone increasing on $[0, \pi)$ onto $[0, \infty)$. On each interval $(m\pi, (m+1)\pi)$, $m \in \mathbb{N}$, μ has an unique critical point b_m , which is a minimum. For $a \geq 0$ and $m \in \mathbb{N}_0$*

$$(2.28) \quad \begin{aligned} (2m+1)\pi &< \tilde{b}_{2m+1} \leq b_{2m+1} \leq 2\hat{b}_m < (2m+2)\pi, \\ (2m+2)\pi &< b_{2m+2} \leq \tilde{b}_{2m+2} < (2m+3)\pi; \end{aligned}$$

and for $a < 0$ and $m \in \mathbb{N}$

$$(2.29) \quad \begin{aligned} 2m\pi &< \tilde{b}_{2m} \leq b_{2m} \leq 2\tilde{b}_m < (2m+1)\pi, \\ (2m+1)\pi &< b_{2m+1} \leq \tilde{b}_{2m+1} < (2m+2)\pi. \end{aligned}$$

On each interval $(m\pi, (m+1)\pi)$, $m \in \mathbb{N}$, μ decreases strictly from $+\infty$ to $\mu(b_m)$ and then increases strictly to $+\infty$. Moreover for all $m \in \mathbb{N}$,

$$(2.30) \quad \mu(b_m) \geq \frac{m-1}{2}\pi,$$

$$(2.31) \quad l(b_m) - b_m \mu(b_m) = 1 - a\delta(b_m),$$

where

$$(2.32) \quad \delta(b) := \cos b + \frac{b}{2} \sin b.$$

Proof. It is easy to see that $m\pi < \tilde{b}_m < m\pi + \frac{\pi}{2} < \hat{b}_m < (m+1)\pi$, see figure 2.




FIGURE 2. The graphs of the functions $\tan$ (black) and $-\cot$ (gray) and the line $x \mapsto x$.

Since

$$\begin{aligned} 1 - b \cot b &= 1 + \frac{b}{2} \tan \frac{b}{2} - \frac{b}{2} \cot \frac{b}{2} \\ &\geq \begin{cases} 1 + \frac{b}{2} \tan \frac{b}{2} & b \in ((2m+1)\pi, (2m+2)\pi) \\ 1 - \frac{b}{2} \cot \frac{b}{2} & b \in (2m\pi, (2m+1)\pi) \end{cases} \end{aligned}$$




FIGURE 3. The graphs of the functions $x \mapsto 1 + \frac{x}{2} \tan \frac{x}{2}$ (black) and $x \mapsto 1 - x \cot x$ (gray).

and $\tilde{b}_{2m+1}, 2\hat{b}_m \in ((2m+1)\pi, (2m+2)\pi)$ it follows that $\tilde{b}_{2m+1} < 2\hat{b}_m$; also, since $\tilde{b}_{2m}, 2\tilde{b}_m \in (2m\pi, (2m+1)\pi)$, it follows that $\tilde{b}_{2m} < 2\tilde{b}_m$.

If $a \geq 0$, μ' is a convex combination of $\tilde{\mu}'$ and $\frac{1}{2}\hat{\mu}'(\frac{b}{2})$. On the interval $((2m+1)\pi, (2m+2)\pi)$ both functions are monotone increasing with zero $\tilde{b}_{2m+1}$ of $\tilde{\mu}'$ lying left of $2\hat{b}_m$, so we have

$$\mu'(\tilde{b}_{2m+1}) = \frac{a}{2}\hat{\mu}'\left(\frac{\tilde{b}_{2m+1}}{2}\right) \leq 0 \text{ and } \mu'(2\hat{b}_m) = (1-a)\tilde{\mu}'(2\hat{b}_m) > 0.$$

Hence μ' has a zero b_m in between: $\tilde{b}_{2m+1} \leq b_m < 2\hat{b}_m$. Due to the fact that μ is strictly convex, this critical point is unique in the interval $((2m+1)\pi, (2m+2)\pi)$, and also a minimum.

In the interval $((2m+2)\pi, (2m+3)\pi)$ still both functions are monotone increasing, but only $\tilde{\mu}'$ has a zero at $\tilde{b}_{2m+2}$. $\frac{1}{2}\hat{\mu}'(\frac{b}{2})$ is positive on this interval and since $\tilde{\mu}'(b)$ tends to $-\infty$ as $b \xrightarrow{b > (2m+2)\pi} (2m+2)\pi$, there is a zero b_{2m+2} of μ' in this interval $((2m+2)\pi, (2m+3)\pi)$ with $(2m+2)\pi \leq b_{2m+2} < \tilde{b}_{2m+2}$. This critical point is unique and a minimum, since μ is convex.

The case $a < 0$ is similar, if we use $\frac{1}{2}\tilde{\mu}(\frac{b}{2})$ instead of $\frac{1}{2}\hat{\mu}(\frac{b}{2})$ and the fact that in the interval $(2m\pi, (2m+1)\pi)$: $\tilde{b}_{2m} < 2\tilde{b}_m$ are zeros of these two functions and $\frac{1}{2}\tilde{\mu}(\frac{b}{2})$ has no zero in $((2m+1)\pi, (2m+2)\pi)$.

To obtain the lower bound (2.30) on μ , we consider the following cases:

1. case $a \geq 0$ and m even:

Since $\frac{b_m}{2} \in ((\frac{m-2}{2} + \frac{1}{2})\pi, (\frac{m}{2} + \frac{1}{2})\pi)$, we have

$$\begin{aligned} \mu(b_m) &= (1-a)\tilde{\mu}(b_m) + a\hat{\mu}\left(\frac{b_m}{2}\right) \\ &\geq (1-a)m\pi + a\frac{m-1}{2}\pi \\ &\geq \frac{m-1}{2}\pi. \end{aligned}$$

2. case $a \geq 0$ and m odd:

Since $\frac{b_m}{2} \in ((\frac{m-1}{2} + \frac{1}{2})\pi, \frac{m+1}{2}\pi)$, we have

$$\begin{aligned}\mu(b_m) &\geq (1-a)m\pi + a\frac{m}{2}\pi \\ &\geq \frac{m}{2}\pi > \frac{m-1}{2}\pi.\end{aligned}$$

3. case $a < 0$ and m even:

Since $\frac{b_m}{2} \in (\frac{m}{2}\pi, \frac{m+1}{2}\pi)$, we have

$$\begin{aligned}\mu(b_m) &= (1+a)\tilde{\mu}(b_m) - a\tilde{\mu}\left(\frac{b_m}{2}\right) \\ &\geq (1+a)m\pi - a\frac{m}{2}\pi \\ &\geq \frac{m}{2}\pi > \frac{m-1}{2}\pi.\end{aligned}$$

4. case $a < 0$ and m odd:

Since $\frac{b_m}{2} \in (\frac{m-1}{2}\pi, \frac{m+1}{2}\pi)$, we have

$$\begin{aligned}\mu(b_m) &\geq (1+a)m\pi - a\frac{m-1}{2}\pi \\ &\geq \frac{m-1}{2}\pi.\end{aligned}$$

At last, to establish the equation (2.31), we need some more information of b_m , i.e. the critical points of μ . First observe, that for any $b \in \mathbb{R} \setminus \pi\mathbb{Z}^*$:

$$(2.33) \quad b\mu(b) + b \cot b - \frac{b}{\sin b}a = l(b)$$

and

$$\begin{aligned}\frac{d}{db} \left(b \cot b - \frac{b}{\sin b}a \right) &= \cot b - \frac{b}{\sin^2 b} - a \frac{1 - b \cot b}{\sin b} \\ &= -\mu(b).\end{aligned}$$

So

$$(2.34) \quad \begin{aligned}l'(b) &= \mu(b) + b\mu'(b) - \mu(b) \\ &= b\mu'(b).\end{aligned}$$

To find the critical points of μ , we differentiate:

$$\begin{aligned}\mu'(b) &= 2 \frac{1 - b \cot b}{\sin^2 b} + a \frac{\frac{b}{\sin b} - \cos b - \cos b(1 - b \cot b)}{\sin^2 b} \\ &= 2 \frac{1 - b \cot b}{\sin^2 b} + a \frac{b \sin b - 2 \cos b(1 - b \cot b)}{\sin^2 b}.\end{aligned}$$

So

$$(2.35) \quad \mu'(b) = 0 \Leftrightarrow a = \frac{1 - b \cot b}{\cos b(1 - b \cot b) - \frac{b}{2} \sin b}.$$

From this and (2.33) we get:

$$\begin{aligned}
l(b_m) - b_m \mu(b_m) &= b_m \cot b_m - \frac{b_m}{\sin b_m} a \\
&= 1 - (1 - b_m \cot b_m) - \frac{b_m}{\sin b_m} a \\
&= 1 - a \left(\cos b_m (1 - b_m \cot b_m) - \frac{b_m}{2} \sin b_m + \frac{b_m}{\sin b_m} \right) \\
&= 1 - a \left(\cos b_m (1 - b_m \cot b_m) + \frac{b_m}{2} \sin b_m + b_m \cos b_m \cot b_m \right) \\
&= 1 - a \left(\cos b_m + \frac{b_m}{2} \sin b_m \right) \\
&= 1 - a \delta(b_m).
\end{aligned}$$

□

For the proof of the next theorem we need another lemma:

Lemma 2.6. *On each interval $(m\pi, (m+1)\pi)$, $m \in \mathbb{N}$ δ has a unique zero, and a unique critical point at $\tilde{b}_m$, which is a maximum with $\delta(\tilde{b}_m) > 0$, if m is even and a minimum with $\delta(\tilde{b}_m) < 0$, if m is odd.*




FIGURE 4. The function δ .

Proof. Since

$$\begin{aligned}
\delta(b) = 0 &\Leftrightarrow \cos b + \frac{b}{2} \sin b = 0 \\
&\Leftrightarrow 2 + b \tan b = 0,
\end{aligned}$$

there is a unique zero in each interval $(m\pi, (m+1)\pi)$, $m \in \mathbb{N}_0$.

The critical points of δ are given by

$$\begin{aligned}\delta'(b) = 0 &\Leftrightarrow -\sin b + \frac{1}{2}\sin b + \frac{b}{2}\cos b = 0 \\ &\Leftrightarrow b \cot b - 1 = 0 \\ &\Leftrightarrow b = \tilde{b}_m.\end{aligned}$$

Since

$$(2.36) \quad \delta''(b) = -\frac{b}{2}\sin b,$$

we see that $\tilde{b}_m$ is a maximum, if m is even and a minimum, if m is odd. $\square$

Theorem 2.3. *Given $(x_1, 0), (x, u) \in \mathbb{R}^n \times \mathbb{R}$ with $x_1 \neq \pm x$ and $u > 0$, there are finitely many geodesics joining these two points. These geodesics are given by (2.6), where b is a solution of*

$$(2.37) \quad \frac{2u}{|x_1|^2 + |x|^2} = \mu(b),$$

and their lengths are strictly increasing with b . Moreover the shortest geodesic joining $(x_1, 0)$ and (x, u) is given by the unique solution $b \in (0, \pi)$ of (2.37) in the interval $(0, \pi)$. With this solution b the Carnot-Carathéodory distance is

$$\begin{aligned}(2.38) \quad d_{CC}((x_1, 0), (x, u)) &= L(\gamma^b) \\ &= \frac{b}{\sin b} \sqrt{|x_1|^2 + |x|^2 - 2x_1 \cdot x \cos b} \\ &= \sqrt{2bu + (|x_1|^2 + |x|^2) b \cot b - 2\frac{b}{\sin b} x_1 \cdot x}.\end{aligned}$$




FIGURE 5. Geodesics joining different points.

Proof. From Lemma 2.1 we have that the geodesics in this case are implicitly given by solutions of the equation $\frac{2u}{|x_1|^2 + |x|^2} = \mu(b)$. The properties of the function μ show us that there is exactly one solution in the interval $(0, \pi)$ and at most two solutions in each interval $(m\pi, (m+1)\pi)$, $m \in \mathbb{N}$. So it remains to prove, that the values of l at these solutions are strictly increasing, i.e. if $s < t$ are two solutions, then $l(s) < l(t)$.

Let $s \in (m\pi, (m+1)\pi)$, $m \in \mathbb{N}$, then we have by (??)

$$\begin{aligned} l(s) &= \int_{b_m}^s l'(b)db + l(b_m) \\ &= \int_{b_m}^s b\mu'(b)db + l(b_m) \\ &= b\mu(b)|_{b_m}^s - \int_{b_m}^s \mu(b)db + l(b_m) \\ &= s\mu(s) - \int_{b_m}^s \mu(b)db + l(b_m) - b_m\mu(b_m) \\ &= s\mu(s) - \int_{b_m}^s \mu(b)db + 1 - a\delta(b_m). \end{aligned}$$

So, if we have two solutions $s \leq t$ of (2.37) in the same interval $(m\pi, (m+1)\pi)$, $m \in \mathbb{N}$, we see that

$$\begin{aligned} l(s) - l(t) &= s\mu(s) - \int_{b_m}^s \mu(b)db - t\mu(t) + \int_{b_m}^t \mu(b)db \\ &= (s - t)\mu(s) + \int_s^t \mu(b)db \\ &\leq 0. \end{aligned}$$

Now let $s_m, t_m \in (m\pi, (m+1)\pi)$, $s_m \leq t_m$, $m \in \mathbb{N}$ be the solutions of (2.37). To compare the corresponding “lengths” l at t_m and s_{m+1} , notice that

$$\begin{aligned} (2.39) \quad l(t_m) - l(s_{m+1}) &= -(s_{m+1} - t_m)\mu(t_m) - \int_{b_m}^{t_m} \mu(b)db - \int_{s_{m+1}}^{b_{m+1}} \mu(b)db - (\delta(b_m) - \delta(b_{m+1}))a. \end{aligned}$$

Now consider the following cases:

1. case $a \geq 0$ and m even:

Then, since $m\pi < b_m \leq \tilde{b}_m$, $\delta(m\pi) = 1$ and $\tilde{b}_m$ is a maximum of δ , it follows that $\delta(b_m) > 1$. And, since $\tilde{b}_{m+1} \leq b_{m+1} < (m+2)\pi$, $\delta((m+2)\pi) = 1$ and $\tilde{b}_{m+1}$ is a minimum of δ , we have $\delta(b_m) < 1$. So $(\delta(b_m) - \delta(b_{m+1}))a \geq 0$, which gives the claim, since $\mu \geq 0$.

2. case $a < 0$ and m odd:

Then, since $m\pi < b_m \leq \tilde{b}_m$, $\delta(m\pi) = -1$ and $\tilde{b}_m$ is a minimum of δ , it follows that $\delta(b_m) < -1$. And, since $\tilde{b}_{m+1} \leq b_{m+1} < (m+2)\pi$, $\delta((m+2)\pi) = -1$ and $\tilde{b}_{m+1}$ is a maximum of δ , we have $\delta(b_m) > -1$. So $(\delta(b_m) - \delta(b_{m+1}))a \geq 0$, which gives the claim, since $\mu \geq 0$.

3. case $a \geq 0$ and m odd:

Here $b_m \leq 2\tilde{b}_{\frac{m-1}{2}} < s_{m+1} \leq b_{m+1}$. Let

$$s := \begin{cases} t_m & , \text{ if } \hat{\mu}\left(\frac{t_m}{2}\right) > \hat{\mu}\left(\frac{s_{m+1}}{2}\right) \\ s_{m+1} & , \text{ if } \hat{\mu}\left(\frac{t_m}{2}\right) \leq \hat{\mu}\left(\frac{s_{m+1}}{2}\right) \end{cases}.$$

Then, since $2\hat{b}_{\frac{m-1}{2}}$ is a minimum of $\hat{\mu}\left(\frac{b}{2}\right)$ on $(m\pi, (m+2)\pi)$ and monotone on the left and right of $2\hat{b}_{\frac{m-1}{2}}$: $\max_{b \in [t_m, s_{m+1}]} \hat{\mu}\left(\frac{b}{2}\right) = \hat{\mu}\left(\frac{s}{2}\right)$ and therefore

$$\begin{aligned} -(s_{m+1} - t_m)\mu(t_m) &= -(s_{m+1} - t_m)\mu(s) \\ &= -(s_{m+1} - t_m)\left((1-a)\tilde{\mu}(s) + a\hat{\mu}\left(\frac{s}{2}\right)\right) \\ &\leq -(s_{m+1} - t_m)a\hat{\mu}\left(\frac{s}{2}\right) \\ &\leq -a \int_{t_m}^{s_{m+1}} \hat{\mu}\left(\frac{b}{2}\right) db. \end{aligned}$$

Notice that $\hat{\mu}\left(\frac{b}{2}\right)$ has no singularity at $(m+1)\pi$. Now (2.39) becomes, since $\mu(b) \geq a\hat{\mu}\left(\frac{b}{2}\right)$:

$$\begin{aligned} l(t_m) - l(s_{m+1}) &\leq -a \int_{b_m}^{b_{m+1}} \hat{\mu}\left(\frac{b}{2}\right) db - (\delta(b_m) - \delta(b_{m+1}))a \\ &= -a \int_{b_m}^{b_{m+1}} \left(\frac{b/2}{\cos^2 \frac{b}{2}} + \tan \frac{b}{2}\right) db - (\delta(b_m) - \delta(b_{m+1}))a \\ &= -ab \tan \frac{b}{2} \Big|_{b_m}^{b_{m+1}} - (\delta(b_m) - \delta(b_{m+1}))a \\ &= a \left(\delta(b_{m+1}) - b_{m+1} \tan \frac{b_{m+1}}{2} - \left(\delta(b_m) - b_m \tan \frac{b_m}{2} \right) \right). \end{aligned}$$

Further

$$\begin{aligned} \delta(b) - b \tan \frac{b}{2} &= \cos b + \frac{b}{2} \sin b - b \tan \frac{b}{2} \\ &= b \sin \frac{b}{2} \left(\cos \frac{b}{2} - \frac{1}{\cos \frac{b}{2}} \right) - 2 \sin^2 \frac{b}{2} + 1 \\ &= - \left(b \sin^2 \frac{b}{2} \tan \frac{b}{2} + 2 \sin^2 \frac{b}{2} \right) + 1 \\ &= -2 \sin^2 \frac{b}{2} \left(1 + \frac{b}{2} \tan \frac{b}{2} \right) + 1. \end{aligned}$$

Since $b_m \leq 2\hat{b}_{\frac{m-1}{2}} \leq b_{m+1}$ and $1 + \hat{b}_{\frac{m-1}{2}} \tan \hat{b}_{\frac{m-1}{2}} = 0$ and $b \mapsto 1 + \frac{b}{2} \tan \frac{b}{2}$ is increasing on $(m\pi, (m+2)\pi)$, we conclude:

$$\begin{aligned} l(t_m) - l(s_{m+1}) &\leq 2a \left(\sin^2 \frac{b_m}{2} \left(1 + \frac{b_m}{2} \tan \frac{b_m}{2} \right) - \sin^2 \frac{b_{m+1}}{2} \left(1 + \frac{b_{m+1}}{2} \tan \frac{b_{m+1}}{2} \right) \right) \\ &\leq 0. \end{aligned}$$

4. case $a < 0$ and m even:

This is very similar to the third case, if we use $\tilde{\mu}$. Here $b_m \leq 2\tilde{b}_{\frac{m}{2}} < s_{m+1} \leq b_{m+1}$. Let

$$s := \begin{cases} t_m & , \text{ if } \tilde{\mu}\left(\frac{t_m}{2}\right) > \tilde{\mu}\left(\frac{s_{m+1}}{2}\right) \\ s_{m+1} & , \text{ if } \tilde{\mu}\left(\frac{t_m}{2}\right) \leq \tilde{\mu}\left(\frac{s_{m+1}}{2}\right) \end{cases}.$$

Then, since $2\tilde{b}_{\frac{m}{2}}$ is a minimum of $\tilde{\mu}\left(\frac{b}{2}\right)$ on $(m\pi, (m+2)\pi)$ and monotone on the left and right of $2\tilde{b}_{\frac{m}{2}}$: $\max_{b \in [t_m, s_{m+1}]} \tilde{\mu}\left(\frac{b}{2}\right) = \tilde{\mu}\left(\frac{s}{2}\right)$ and therefore

$$\begin{aligned} -(s_{m+1} - t_m)\mu(t_m) &= -(s_{m+1} - t_m)\mu(s) \\ &= -(s_{m+1} - t_m)\left((1+a)\tilde{\mu}(s) - a\tilde{\mu}\left(\frac{s}{2}\right)\right) \\ &\leq (s_{m+1} - t_m)a\tilde{\mu}\left(\frac{s}{2}\right) \\ &\leq a \int_{t_m}^{s_{m+1}} \tilde{\mu}\left(\frac{b}{2}\right) db. \end{aligned}$$

Notice that $\tilde{\mu}\left(\frac{b}{2}\right)$ has no singularity at $(m+1)\pi$. Now (2.39) becomes, since $\mu(b) \geq -a\tilde{\mu}(b/2)$:

$$\begin{aligned} l(t_m) - l(s_{m+1}) &\leq a \left(\int_{b_m}^{b_{m+1}} \tilde{\mu}\left(\frac{b}{2}\right) db - \delta(b_m) + \delta(b_{m+1}) \right) \\ &= a \left(\int_{b_m}^{b_{m+1}} \left(\frac{b/2}{\sin^2 \frac{b}{2}} - \cot \frac{b}{2} \right) db - \delta(b_m) + \delta(b_{m+1}) \right) \\ &= a \left(-b \cot \frac{b}{2} \Big|_{b_m}^{b_{m+1}} - \delta(b_m) + \delta(b_{m+1}) \right) \\ &= a \left(b_m \cot \frac{b_m}{2} - \delta(b_m) - (b_{m+1} \cot \frac{b_{m+1}}{2} - \delta(b_{m+1})) \right). \end{aligned}$$

Further

$$\begin{aligned} b \cot \frac{b}{2} - \delta(b) &= -\cos b - \frac{b}{2} \sin b + b \cot \frac{b}{2} \\ &= -b \cos \frac{b}{2} \left(\sin \frac{b}{2} - \frac{1}{\sin \frac{b}{2}} \right) - 2 \cos^2 \frac{b}{2} + 1 \\ &= -2 \cos^2 \frac{b}{2} \left(1 - \frac{b}{2} \cot \frac{b}{2} \right) + 1. \end{aligned}$$

Since $b_m \leq 2\tilde{b}_{\frac{m}{2}} \leq b_{m+1}$ and $1 - \tilde{b}_{\frac{m}{2}} \cot \tilde{b}_{\frac{m}{2}} = 0$ and $b \mapsto 1 - \frac{b}{2} \cot \frac{b}{2}$ is increasing on $(m\pi, (m+2)\pi)$, we conclude:

$$\begin{aligned} l(t_m) - l(s_{m+1}) &\leq -2a \left(\cos^2 \frac{b_m}{2} \left(1 - \frac{b_m}{2} \cot \frac{b_m}{2} \right) - \cos^2 \frac{b_{m+1}}{2} \left(1 - \frac{b_{m+1}}{2} \cot \frac{b_{m+1}}{2} \right) \right) \\ &\leq 0. \end{aligned}$$

If $t_0 \in (0, \pi)$ denotes the solution of (2.37), it remains to show that $l(t_0) < l(s_1)$. The above calculations can be extended to this, if one sets $b_0 = 0$. In detail we have

$$\begin{aligned} l(t_0) &= \int_0^{t_0} l'(b) db + l(0) \\ &= t_0 \mu(t_0) - \int_0^{t_0} \mu(b) db + 1 - a, \end{aligned}$$

so that

$$l(t_0) - l(s_1) = -(s_1 - t_0)\mu(s_1) - \int_0^{t_0} \mu(b) db - \int_{s_1}^{b_1} \mu(b) db - (1 - \delta(b_1))a.$$

Again from the previous lemma one gets $l(t_0) < l(s_1)$ if $a \geq 0$, since $1 - \delta(b_1) > 0$ and $\mu > 0$. If $a < 0$, then since $\tilde{\mu}$ is increasing on $[0, \pi)$:

$$\begin{aligned} -(s_1 - t_0)\mu(s_1) &= -(s_1 - t_0) \left((1 + a)\tilde{\mu}(s_1) - a\tilde{\mu}\left(\frac{s_1}{2}\right) \right) \\ &\leq (s_1 - t_0)a\tilde{\mu}\left(\frac{s_1}{2}\right) \\ &\leq a \int_{t_0}^{s_1} \tilde{\mu}\left(\frac{b}{2}\right) db. \end{aligned}$$

So we get

$$\begin{aligned} l(t_0) - l(s_1) &\leq a \int_0^{b_1} \tilde{\mu}\left(\frac{b}{2}\right) db - (1 - \delta(b_1))a \\ &= a \left(-b \cot \frac{b}{2} \Big|_0^{b_1} - 1 + \delta(b_1) \right) \\ &= a \left(\delta(b_1) - b_1 \cot \frac{b_1}{2} + 1 \right) \\ &= 2a \cos^2 \frac{b_1}{2} \left(1 - \frac{b_1}{2} \cot \frac{b_1}{2} \right) \\ &\leq 0, \end{aligned}$$

since $1 - \frac{b}{2} \cot \frac{b}{2} \geq 0$ on $[0, 2\pi)$. □

Theorem 2.4. *Given $(x, 0), (x, u) \in \mathbb{R}^n \times \mathbb{R}$, $x \neq 0$, $u > 0$, there are finitely many geodesics connecting $(x, 0)$, (x, u) . Namely:*

1. *For any solution b of*

$$(2.40) \quad \frac{u}{|x|^2} = \hat{\mu}\left(\frac{b}{2}\right)$$

there is a geodesic γ^b given by (2.6) with square of length

$$(2.41) \quad L(\gamma^b)^2 = \frac{b^2}{\cos^2 \frac{b}{2}} |x|^2 = 2bu - 2|x|^2 b \tan \frac{b}{2},$$

which strictly increases with b .

2. *If $u \geq |x|^2 \pi$, then for each $m \in \mathbb{N}$ with $u \geq m\pi |x|^2$ there are geodesics $\gamma^{b,c}$ given by (2.6) with $b = 2m\pi$, $|c| = 2\sqrt{m\pi u - (m\pi)^2 |x|^2}$, with lengths*

$$(2.42) \quad L(\gamma^{b,c}) = 2\sqrt{m\pi u}.$$

Moreover the shortest geodesic and therefore the Carnot-Carathéodory distance is given by the unique solution b of (2.40) in the interval $(0, \pi)$, with this b we have

$$(2.43) \quad d_{CC}((x, 0), (x, u)) = \frac{b}{\cos \frac{b}{2}} |x| = \sqrt{2bu - 2|x|^2 b \tan \frac{b}{2}}.$$

Proof. Notice that in this case $a = 1$. We will consider two cases:

1. **case $b \in \pi\mathbb{N}$:**

Then since $\gamma_{(1)}^{b,c}(1) = x$, there exists $m \in \mathbb{N}$ with $b = 2m\pi$. Then $\gamma_{(2)}^{b,c}(1) = u$ is equivalent to

$$u = \left(|c|^2 + (2m\pi)^2 |x|^2 \right) \frac{1}{4m\pi}.$$




FIGURE 6. Geodesics joining $(1,0)$ and $(1,6)$ with different $b \notin \mathbb{Z}^*$ vs. the two geodesics with $b = 2\pi$ and $c = \pm\sqrt{2bu - x_1^2 b^2}$.

So if $u \geq m\pi|x|^2$, there is a geodesic given by (2.6) with

$$b = 2m\pi, \quad |c| = 2\sqrt{m\pi u - (m\pi)^2|x|^2},$$

with length:

$$L^2(\gamma^{b,c}) = |c|^2 + |x|^2 b^2 = 4m\pi u$$

This gives the second part of the theorem.

2. case $b \notin \pi\mathbb{N}$:

In this case we have

$$\mu(b) = \hat{\mu}\left(\frac{b}{2}\right).$$

This is very similar to the situation in Theorem 2.3, but much easier, since we are only dealing with $\hat{\mu}$. From Lemma 2.4 we know that the equation

$$\frac{u}{|x|^2} = \hat{\mu}\left(\frac{b}{2}\right)$$

has at most two solutions in an interval $((2m+1)\pi, (2m+3)\pi)$, $m \in \mathbb{N}_0$, each defining a geodesic. If s is a solution of this equation, the square of the length of the corresponding geodesic γ is given by

$$\begin{aligned} L^2(\gamma) &= 2l(s)|x|^2 \\ &= 2\frac{s^2}{\sin^2 s}(1 - \cos s)|x|^2 \\ &= \frac{s^2}{\cos^2 \frac{s}{2}}|x|^2. \end{aligned}$$

If $s \in ((2m+1)\pi, (2m+3)\pi)$, $m \in \mathbb{N}_0$, we have

$$\begin{aligned}
 l(s) &= \int_{2\hat{b}_m}^s l'(b)db + l(2\hat{b}_m) \\
 &= \int_{2\hat{b}_m}^s b\mu'(b)db + l(2\hat{b}_m) \\
 &= \int_{2\hat{b}_m}^s \frac{b}{2}\hat{\mu}'\left(\frac{b}{2}\right)db + l(2\hat{b}_m) \\
 (2.44) \quad &= s\hat{\mu}\left(\frac{s}{2}\right) - 2\hat{b}_m\hat{\mu}(\hat{b}_m) - \int_{2\hat{b}_m}^s \hat{\mu}\left(\frac{b}{2}\right)db + l(2\hat{b}_m)
 \end{aligned}$$

But now

$$\begin{aligned}
 l(2\hat{b}_m) - 2\hat{b}_m\hat{\mu}(\hat{b}_m) &= \frac{2\hat{b}_m^2}{\cos^2 \hat{b}_m} - 2\hat{b}_m \left(\frac{\hat{b}_m}{\cos^2 \hat{b}_m} + \tan \hat{b}_m \right) \\
 &= -2\hat{b}_m \tan \hat{b}_m \\
 (2.45) \quad &= 2.
 \end{aligned}$$

First let $s \leq t$ be two solutions in an interval $((2m+1)\pi, (2m+3)\pi)$, $m \in \mathbb{N}_0$. Then by (??) and (??)

$$\begin{aligned}
 l(s) - l(t) &= s\hat{\mu}\left(\frac{s}{2}\right) - \int_{2\hat{b}_m}^s \hat{\mu}\left(\frac{b}{2}\right)db - t\hat{\mu}\left(\frac{t}{2}\right) + \int_{2\hat{b}_m}^t \hat{\mu}\left(\frac{b}{2}\right)db \\
 &= -(t-s)\hat{\mu}\left(\frac{s}{2}\right) + \int_s^t \hat{\mu}\left(\frac{b}{2}\right)db \\
 &\leq 0.
 \end{aligned}$$

Now let $t_m \in ((2m+1)\pi, (2m+3)\pi)$ and $s_{m+1} \in ((2m+3)\pi, (2m+5)\pi)$ be two solutions, where t_m is the rightmost and s_{m+1} is the leftmost solution in the corresponding interval. Then by (??) and (??)

$$\begin{aligned}
 l(t_m) - l(s_{m+1}) &= t_m\hat{\mu}\left(\frac{t_m}{2}\right) - \int_{2\hat{b}_m}^{t_m} \hat{\mu}\left(\frac{b}{2}\right)db - s_{m+1}\hat{\mu}\left(\frac{s_{m+1}}{2}\right) + \int_{2\hat{b}_{m+1}}^{s_{m+1}} \hat{\mu}\left(\frac{b}{2}\right)db \\
 &= -(s_{m+1} - t_m)\hat{\mu}\left(\frac{t_m}{2}\right) - \int_{2\hat{b}_m}^{t_m} \hat{\mu}\left(\frac{b}{2}\right)db - \int_{s_{m+1}}^{2\hat{b}_{m+1}} \hat{\mu}\left(\frac{b}{2}\right)db \\
 &\leq 0,
 \end{aligned}$$

since $2\hat{b}_m \leq t_m$ and $s_{m+1} \leq 2\hat{b}_{m+1}$.

If $t \in (0, \pi)$ is the solution of (2.40), we have to check, that $l(t) < l(s_0)$. As above, we have

$$\begin{aligned}
 l(t) &= \int_0^t l'(b)db + l(0) \\
 &= t\hat{\mu}\left(\frac{t}{2}\right) - \int_0^t \hat{\mu}\left(\frac{b}{2}\right)db + 2,
 \end{aligned}$$

so that

$$\begin{aligned}
 l(t) - l(s_0) &= t\hat{\mu}\left(\frac{t}{2}\right) - \int_0^t \hat{\mu}\left(\frac{b}{2}\right)db - s_0\hat{\mu}\left(\frac{s_0}{2}\right) + \int_{2\hat{b}_0}^{s_0} \hat{\mu}\left(\frac{b}{2}\right)db \\
 &= -(s_0 - t)\hat{\mu}\left(\frac{t}{2}\right) - \int_0^t \hat{\mu}\left(\frac{b}{2}\right)db - \int_{s_0}^{2\hat{b}_0} \hat{\mu}\left(\frac{b}{2}\right)db \\
 &\leq 0.
 \end{aligned}$$

At last we have to compare the length of the shortest geodesic of case 1 (if there is one) with that of case 2. Let $b \in (0, \pi)$ be the solution of (2.40), then since $\tan \frac{b}{2} \geq 0$:

$$\begin{aligned} L^2(\gamma^b) &= 2bu - 2|x|^2 b \tan \frac{b}{2} \\ &\leq 2\pi u \\ &\leq L^2(\gamma^{2\pi, c}). \end{aligned}$$

□

Theorem 2.5. *Given $(x, 0), (-x, u) \in \mathbb{R}^n \times \mathbb{R}$, $x \neq 0$, $u > 0$, there are finitely many geodesics connecting $(x, 0)$, $(-x, u)$. Namely:*

1. *For any solution b of*

$$(2.46) \quad \frac{u}{|x|^2} = \tilde{\mu} \left(\frac{b}{2} \right)$$

there is a geodesic γ^b given by (2.6) with square of length

$$(2.47) \quad L(\gamma^b)^2 = \frac{b^2}{\sin^2 \frac{b}{2}} |x|^2 = 2bu + |x|^2 b \cot \frac{b}{2},$$

which strictly increases with b .

2. *If $2u \geq |x|^2 \pi$, then for each $m \in \mathbb{N}_0$ with $2u \geq (2m+1)\pi |x|^2$ there are geodesics $\gamma^{b,c}$ given by (2.6) with $b = (2m+1)\pi$, $|c| = \sqrt{2(2m+1)\pi u - (2m+1)^2 \pi^2 |x|^2}$ and lengths*

$$(2.48) \quad L(\gamma^{b,c}) = \sqrt{2(2m+1)\pi u}.$$

Moreover the shortest geodesic and therefore the Carnot-Carathéodory distance is given by the unique solution b of (2.46) in the interval $(0, \pi)$, if $2u < |x|^2 \pi$. With this b we have:

$$(2.49) \quad d_{CC}((x, 0), (-x, u)) = \frac{b}{\sin \frac{b}{2}} |x| = \sqrt{2bu + |x|^2 b \cot \frac{b}{2}}.$$

And, if $2u \geq |x|^2 \pi$, then the Carnot-Carathéodory distance is given by

$$d_{CC}((x, 0), (-x, u)) = \sqrt{2\pi u}.$$

Proof. Notice that in this case $a = -1$. The first part is the same proof as above, if one replaces $\hat{\mu}$, $\hat{b}$ with $\tilde{\mu}$, $\tilde{b}$. In particular:

1. **case $b \in \pi\mathbb{N}$:**

Then since $\gamma_{(1)}(1) = x$, there exists $m \in \mathbb{N}_0$ with $b = (2m+1)\pi$. Then $\gamma_{(2)}(1) = u$ is equivalent to

$$u = \left(|c|^2 + (2m+1)^2 \pi^2 |x|^2 \right) \frac{1}{2(2m+1)\pi}.$$

So if $2u \geq (2m+1)\pi |x|^2$, there are geodesics $\gamma^{b,c}$ given by (2.6) with $b = (2m+1)\pi$, $|c| = \sqrt{2(2m+1)\pi u - (2m+1)^2 \pi^2 |x|^2}$, with length

$$L^2(\gamma^{b,c}) = |c|^2 + |x|^2 b^2 = 2(2m+1)\pi u$$

This gives the second part of the theorem.




FIGURE 7. Geodesics joining $(1, 0)$ and $(-1, 5)$ with different $b \notin \mathbb{Z}^*$ vs. two geodesics with $b = \pi$ and $c = \pm \sqrt{2bu - x_1^2 b^2}$.

2. case $b \notin \pi\mathbb{N}$:

In this case we have:

$$\mu(b) = \tilde{\mu}\left(\frac{b}{2}\right).$$

This is again very similar to the situation in Theorem 2.3, but a lot simpler, since we are only dealing with $\tilde{\mu}$. From Lemma 2.3 we know that the equation

$$(2.50) \quad \frac{u}{|x|^2} = \tilde{\mu}\left(\frac{b}{2}\right)$$

has at most two solutions in an interval $(2m\pi, (2m+2)\pi)$, $m \in \mathbb{N}$, each defining an geodesic. If s is a solution of this equation, the square of the length of the corresponding geodesic γ is given by

$$\begin{aligned} L^2(\gamma) &= 2l(s)|x|^2 \\ &= 2 \frac{s^2}{\sin^2 s} (1 + \cos s) |x|^2 \\ &= \frac{s^2}{\sin^2 \frac{s}{2}} |x|^2. \end{aligned}$$

If $s \in (2m\pi, (2m+2)\pi)$, $m \in \mathbb{N}$, we have:

$$\begin{aligned} l(s) &= \int_{2\tilde{b}_m}^s l'(b)db + l(2\tilde{b}_m) \\ &= \int_{2\tilde{b}_m}^s b\mu'(b)db + l(2\tilde{b}_m) \\ &= \int_{2\tilde{b}_m}^s \frac{b}{2} \tilde{\mu}'\left(\frac{b}{2}\right) db + l(2\tilde{b}_m) \\ &= s\tilde{\mu}\left(\frac{s}{2}\right) - 2\tilde{b}_m\tilde{\mu}\left(\tilde{b}_m\right) - \int_{2\tilde{b}_m}^s \tilde{\mu}\left(\frac{b}{2}\right) db + l(2\tilde{b}_m) \end{aligned}$$

But now

$$\begin{aligned} l(2\tilde{b}_m) - 2\tilde{b}_m\tilde{\mu}(\tilde{b}_m) &= \frac{2\tilde{b}_m^2}{\sin^2 \tilde{b}_m} - 2\tilde{b}_m \left(\frac{\tilde{b}_m}{\sin^2 \tilde{b}_m} - \cot \tilde{b}_m \right) \\ &= 2\tilde{b}_m \cot \tilde{b}_m \\ &= 2. \end{aligned}$$

First let $s \leq t$ be two solutions of (2.50) in an interval $(2m\pi, (2m+2)\pi)$, $m \in \mathbb{N}$; then

$$\begin{aligned} l(s) - l(t) &= s\tilde{\mu}\left(\frac{s}{2}\right) - \int_{2\tilde{b}_m}^s \tilde{\mu}\left(\frac{b}{2}\right) db - t\tilde{\mu}\left(\frac{t}{2}\right) + \int_{2\tilde{b}_m}^t \tilde{\mu}\left(\frac{b}{2}\right) db \\ &= -(t-s)\tilde{\mu}\left(\frac{s}{2}\right) + \int_s^t \tilde{\mu}\left(\frac{b}{2}\right) db \\ &\leq 0. \end{aligned}$$

Now let $t_m \in (2m\pi, (2m+2)\pi)$ and $s_{m+1} \in ((2m+2)\pi, (2m+4)\pi)$ be two solutions, then

$$\begin{aligned} l(t_m) - l(s_{m+1}) &= t_m\tilde{\mu}\left(\frac{t_m}{2}\right) - \int_{2\tilde{b}_m}^{t_m} \tilde{\mu}\left(\frac{b}{2}\right) db - s_{m+1}\tilde{\mu}\left(\frac{s_{m+1}}{2}\right) + \int_{2\tilde{b}_{m+1}}^{s_{m+1}} \tilde{\mu}\left(\frac{b}{2}\right) db \\ &= -(s_{m+1} - t_m)\tilde{\mu}\left(\frac{t_m}{2}\right) - \int_{2\tilde{b}_m}^{t_m} \tilde{\mu}\left(\frac{b}{2}\right) db - \int_{s_{m+1}}^{2\tilde{b}_{m+1}} \tilde{\mu}\left(\frac{b}{2}\right) db \\ &\leq 0, \end{aligned}$$

since $2\tilde{b}_m \leq t_m$ and $s_{m+1} \leq 2\tilde{b}_{m+1}$.

If $t_0 \in (0, 2\pi)$ is the solution in the interval $(0, 2\pi)$, we have to show that $l(t_0) < l(s_1)$. But the above calculations show that

$$\begin{aligned} l(t_0) &= \int_0^{t_0} l'(b) db + l(0) \\ &= t_0\tilde{\mu}\left(\frac{t_0}{2}\right) - \int_0^{t_0} \tilde{\mu}\left(\frac{b}{2}\right) db + 2. \end{aligned}$$

So we get

$$\begin{aligned} l(t_0) - l(s_1) &= t_0\tilde{\mu}\left(\frac{t_0}{2}\right) - \int_0^{t_0} \tilde{\mu}\left(\frac{b}{2}\right) db - s_1\tilde{\mu}\left(\frac{s_1}{2}\right) + \int_{2\tilde{b}_1}^{s_1} \tilde{\mu}\left(\frac{b}{2}\right) db \\ &= -(s_1 - t_0)\tilde{\mu}\left(\frac{t_0}{2}\right) - \int_0^{t_0} \tilde{\mu}\left(\frac{b}{2}\right) db - \int_{s_1}^{2\tilde{b}_1} \tilde{\mu}\left(\frac{b}{2}\right) db \\ &< 0. \end{aligned}$$

At last we have to compare the length of the shortest geodesic $\gamma^{\pi, c}$, $|c| = \sqrt{2\pi u - \pi^2 |x|^2}$ of case 1 (if there is one) with that of case 2. Let $b \in (0, 2\pi)$ be the solution of (2.46). If $b < \pi$, then, since $\tilde{\mu}$ is monotone increasing on $(0, \pi)$, we get:

$$\begin{aligned} \frac{u}{|x|^2} &= \tilde{\mu}\left(\frac{b}{2}\right) \\ &< \tilde{\mu}\left(\frac{\pi}{2}\right) \\ &= \frac{\pi}{2}. \end{aligned}$$

But this means, that $2u < \pi |x|^2$ and therefore there is no extra geodesic. If now $b \geq \pi$, we have to show that $L^2(\gamma^b) \geq 2\pi u$. We have

$$\begin{aligned} \frac{u}{|x|^2} &= \tilde{\mu}\left(\frac{b}{2}\right) \\ &= \frac{\frac{b}{2}}{\sin^2 \frac{b}{2}} - \cot \frac{b}{2} \\ &= \cot \frac{b}{2} \left(\frac{b}{2 \sin \frac{b}{2} \cos \frac{b}{2}} - 1 \right) \\ &= \frac{b - \sin b}{\sin b} \cot \frac{b}{2}, \end{aligned}$$

so that

$$\begin{aligned} L^2(\gamma^b) &= 2bu + 2|x|^2 b \cot \frac{b}{2} \\ &= 2bu \left(1 + \frac{\sin b}{b - \sin b} \right) \\ &= 2u \frac{b^2}{b - \sin b} \\ &\geq 2\pi u. \end{aligned}$$

The last inequality holds since $f(b) := \frac{b^2}{b - \sin b} \geq \pi$, for all $b \in [\pi, 2\pi]$. To see this, observe, that $f(\pi) = \pi$ and f is monotone increasing:

$$\begin{aligned} f'(b) &= \frac{2b(b - \sin b) - b^2(1 - \cos b)}{(b - \sin b)^2} \\ &= \frac{b^2(1 + \cos b) - 2b \sin b}{(b - \sin b)^2} \\ &\geq 0. \end{aligned}$$

This completes the proof, since the geodesics $\gamma^{b,c}$ in case 2 (with parameter $b = \pi$, $|c| = \sqrt{2\pi u - \pi^2 x^2}$) are shorter. $\square$

To close this section, we give a summarizing theorem about our results of this section:

Theorem 2.6. *Let $(x_1, u_1), (x, u) \in \mathbb{R}^{n+1}$. If $x_1 = -x$ and $2|u - u_1| \geq \pi |x|^2$, then the Carnot-Carathéodory distance is given by*

$$(2.51) \quad d_{CC}((x_1, u_1), (x, u)) = \sqrt{2\pi |u - u_1|}.$$

Otherwise the Carnot-Carathéodory distance is given by

$$(2.52a) \quad d_{CC}((x_1, u_1), (x, u)) = \frac{b}{\sin b} \sqrt{|x_1|^2 + |x|^2 - 2x_1 \cdot x \cos b}$$

$$(2.52b) \quad = \sqrt{2b(u - u_1) + (|x_1|^2 + |x|^2)b \cot b - 2x_1 \cdot x \frac{b}{\sin b}}$$

where $b \in (-\pi, \pi)$ is the unique solution of

$$(2.53) \quad \frac{2(u - u_1)}{|x_1|^2 + |x|^2} = \frac{b}{\sin^2 b} - \cot b + 2 \frac{x_1 \cdot x}{|x_1|^2 + |x|^2} \cdot \frac{1 - b \cot b}{\sin b}$$

in the interval $(-\pi, \pi)$.

3. HEAT KERNEL

We now turn our attention to the heat equation, i.e. the Cauchy problem

$$(3.1) \quad \begin{cases} (\partial_t - G)v(t, x, u) = 0 \\ v(0, x, u) = f(x, u) \end{cases}$$

with some function $f \in \mathcal{S}(\mathbb{R}^{n+1})$. Taking the Fourier-transform in the u -variable we see that this is equivalent to

$$(3.2) \quad \begin{cases} (\partial_t - H_\lambda)w(t, x, \lambda) = 0 \\ w(0, x, \lambda) = g(x, \lambda) \end{cases},$$

where $g(x, \lambda) := f(x, \hat{\lambda})$ denotes the partial Fourier-transform of f in the u -variable and $H_\lambda = \Delta_x - |\lambda|^2|x|^2$ denote the Hermite-operator with parameter $\lambda \in \mathbb{R}$, which is the partial Fourier-transform of G . Thanks to the Plancherel formula, any solution w of (3.2) defines a solution v to (3.1) by taking the inverse Fourier-transform, and vice versa.

But the latter equation is well known and we give a brief discussion of this equation. The solution of this equation is given by $e^{tH_\lambda}g$, where e^{tH_λ} is a bounded operator defined by the functional calculus, since H_λ is a self-adjoint operator. By the Schwartz kernel theorem e^{tH_λ} is given by an integral kernel, i.e.

$$(3.3) \quad e^{tH_\lambda}g(x) = \int_{\mathbb{R}^n} k_t^\lambda(x, \xi)g(\xi)d\xi.$$

And, since H_λ is hypoelliptic, we have $k_t^\lambda \in C^\infty(\mathbb{R}^{2n})$.

The Hermite-functions

$$(3.4) \quad h_\alpha(x) := \frac{2^{n/4}}{\sqrt{\alpha!}} \left(\frac{-1}{2\sqrt{\pi}} \right)^{|\alpha|} e^{\pi|x|^2} \left(\frac{\partial}{\partial x} \right)^\alpha e^{-2\pi|x|^2}, \quad x \in \mathbb{R}^n, \alpha \in \mathbb{N}_0^n,$$

which give an orthonormal basis of $L^2(\mathbb{R}^n)$, are eigenfunctions of $H_{2\pi}$:

$$(3.5) \quad H_{2\pi}h_\alpha = -2\pi(2|\alpha| + n)h_\alpha.$$

Since

$$(3.6) \quad (\Delta - \mu^2|x|^2)f(r\cdot) = r^2 \left(\left(\Delta - \left(\frac{\mu}{r^2} \right)^2 |\cdot|^2 \right) f \right)(r\cdot),$$

we can rescale the Hermite functions

$$(3.7) \quad h_\alpha^\lambda(x) := \left(\frac{|\lambda|}{2\pi} \right)^{\frac{n}{4}} h_\alpha \left(\left(\frac{|\lambda|}{2\pi} \right)^{\frac{1}{2}} x \right)$$

to get an orthonormal basis $(h_\alpha^\lambda)_{\alpha \in \mathbb{N}_0^n}$ of eigenfunctions of H_λ :

$$(3.8) \quad H_\lambda h_\alpha^\lambda = -|\lambda|(2|\alpha| + n)h_\alpha^\lambda.$$

Using Mehler's formula, we can calculate the kernel k_t^λ :

$$\begin{aligned}
k_t^\lambda(x, \xi) &= \sum_{|\alpha| \geq 0} e^{tH_\lambda} h_\alpha^\lambda(x) h_\alpha^\lambda(\xi) \\
&= \sum_{|\alpha| \geq 0} e^{-|\lambda|(2|\alpha|+n)t} \left(\frac{|\lambda|}{2\pi} \right)^{\frac{n}{2}} h_\alpha \left(\sqrt{\frac{|\lambda|}{2\pi}} x \right) h_\alpha \left(\sqrt{\frac{|\lambda|}{2\pi}} \xi \right) \\
&= \left(\frac{|\lambda|}{2\pi} \right)^{\frac{n}{2}} e^{-n|\lambda|t} \sum_{|\alpha| \geq 0} e^{-2|\lambda||\alpha|t} h_\alpha \left(\sqrt{\frac{|\lambda|}{2\pi}} x \right) h_\alpha \left(\sqrt{\frac{|\lambda|}{2\pi}} \xi \right) \\
&= \left(\frac{|\lambda|}{2\pi} \right)^{\frac{n}{2}} e^{-n|\lambda|t} \left(\frac{2}{1 - e^{-4|\lambda|t}} \right)^{\frac{n}{2}} \\
&\quad \cdot \exp \left(\frac{-\pi (1 + e^{-4|\lambda|t}) \frac{|x|^2 + |\xi|^2}{2\pi} |\lambda| + 4\pi \frac{x \cdot \xi}{2\pi} |\lambda| e^{-2|\lambda|t}}{1 - e^{-4|\lambda|t}} \right) \\
&= (2\pi)^{-n/2} \left(\frac{|\lambda|}{\sinh(2|\lambda|t)} \right)^{\frac{n}{2}} \\
&\quad \cdot \exp \left(-\frac{|x|^2 + |\xi|^2}{2} |\lambda| \coth(2|\lambda|t) + \frac{|\lambda|}{\sinh(2|\lambda|t)} x \cdot \xi \right) \\
&= (4\pi t)^{-n/2} \left(\frac{2\lambda t}{\sinh(2\lambda t)} \right)^{\frac{n}{2}} \\
&\quad \cdot \exp \left(-\frac{1}{4t} \left(2\lambda t \coth(2\lambda t) (|x|^2 + |\xi|^2) - \frac{4\lambda t}{\sinh(2\lambda t)} x \cdot \xi \right) \right).
\end{aligned}$$

Now, by taking the Fourier inverse

$$\begin{aligned}
(3.9) K_t(x, \xi, u) &= (2\pi)^{-1} \int_{-\infty}^{\infty} k_t^\lambda(x, \xi) e^{i\lambda u} d\lambda \\
&= (2\pi)^{-1} (4\pi t)^{-n/2} \int_{-\infty}^{\infty} \left(\frac{2\lambda t}{\sinh(2\lambda t)} \right)^{\frac{n}{2}} \\
&\quad \cdot \exp \left(-\frac{1}{4t} \left(2\lambda t \coth(2\lambda t) (|x|^2 + |\xi|^2) - \frac{4\lambda t}{\sinh(2\lambda t)} x \cdot \xi \right) \right) e^{i\lambda u} d\lambda \\
&= (4\pi t)^{-n/2-1} \int_{-\infty}^{\infty} \left(\frac{\lambda}{\sinh \lambda} \right)^{\frac{n}{2}} \\
&\quad \exp \left(-\frac{1}{4t} \left(\lambda \coth \lambda (|x|^2 + |\xi|^2) - \frac{2\lambda}{\sinh \lambda} x \cdot \xi - 2i\lambda u \right) \right) d\lambda,
\end{aligned}$$

one gets the heat kernel for the Grušin operator. Now the solution of (3.1) is given by

$$(3.10) \quad v(x, u) = \int_{\mathbb{R}} \int_{\mathbb{R}^n} K_t(x, \xi, u - \lambda) f(\xi, \lambda) d\xi d\lambda.$$

4. ESTIMATES

We may now use the results of section 2 to give some estimates for the heat kernel. We set

$$(4.1) \quad \begin{aligned} h(x, \xi, u) &:= \int_{-\infty}^{\infty} \left(\frac{\lambda}{\sinh \lambda} \right)^{\frac{n}{2}} \exp \left(- \left(\lambda \coth \lambda (|x|^2 + |\xi|^2) - \frac{2\lambda}{\sinh \lambda} x \cdot \xi - 2i\lambda u \right) \right) d\lambda \\ &= \int_{-\infty}^{\infty} \left(\frac{\lambda}{\sinh \lambda} \right)^{\frac{n}{2}} \exp \left(- \left(\lambda \coth \lambda - \frac{\lambda}{\sinh \lambda} a \right) R^2 + 2i\lambda u \right) d\lambda \end{aligned}$$

with

$$R := R(x, \xi) := \sqrt{|x|^2 + |\xi|^2} \text{ and } a := a(x, \xi) := \frac{2x \cdot \xi}{R^2} \in [0, 1].$$

Then

$$(4.2) \quad K_t(x, \xi, u) = (4\pi t)^{-n/2-1} h \left(\frac{x}{2\sqrt{t}}, \frac{\xi}{2\sqrt{t}}, \frac{u}{4t} \right).$$

It is also convenient to set

$$(4.3) \quad V(\lambda) := \left(\frac{\lambda}{\sinh \lambda} \right)^{\frac{n}{2}}$$

$$(4.4) \quad \begin{aligned} \psi(\lambda) &:= \psi(\lambda, a) \\ &:= \lambda \coth \lambda - \frac{\lambda}{\sinh \lambda} a, \quad \lambda \in \mathbb{R}. \end{aligned}$$

One may see that $\psi(ib) = b \cot b - \frac{b}{\sin b} a$, so that if b is a solution of (2.53), then the exponent

$$\left(\lambda \coth \lambda - \frac{\lambda}{\sinh \lambda} a \right) R^2 - 2i\lambda u$$

at $\lambda = ib$ gives exactly the square of the Carnot-Carathéodory distance $d_{CC}((x, 0), (\xi, u))^2$.

So we expect Gaussian-type estimates of the form

$$|h(x, \xi, u)| \leq F(x, \xi, u) e^{-d_{CC}((x, 0), (\xi, u))^2},$$

where $F(x, \xi, u) > 0$ is a function depending on x, ξ, u . It will turn out that F has polynomial growth, more precisely

$$F(x, \xi, u) \lesssim (1 + d_{CC}((x, 0), (\xi, u))^2)^\alpha,$$

where $\alpha = \frac{n}{2} - 1$, if $n > 2$ and $\alpha = 0$ if $n \leq 2$. Compared to the euclidean case, where $\alpha = 0$, one has some additional growth, if the dimension is greater than 2.

The strategy is as follows: Move the line of integration in (4.1) from the real axis to the line $\mathbb{R} + ib$, where b is near to the solution b_0 of (2.53).

Lemma 4.1. *The function*

$$V(\lambda) := \left(\frac{\lambda}{\sinh \lambda} \right)^{\frac{n}{2}},$$

where the square root is the principal branch in $\mathbb{C} \setminus (-\infty, 0]$, is holomorphic in $\{z \in \mathbb{C} : |\Im z| < \pi\}$. And for $\nu + ib \in \mathbb{C}$, $-\pi < b < \pi$, one has

$$(4.5) \quad |V(\nu + ib)| \leq \left(\frac{b}{\sin b} \right)^{\frac{n}{2}},$$

$$(4.6) \quad |V(\nu + ib)| \leq \left(1 + \frac{b^2}{\nu^2} \right)^{\frac{n}{4}} \left(\frac{\nu}{\sinh \nu} \right)^{\frac{n}{2}}.$$

Proof. To show that V is holomorphic, we have to show that

$$\frac{\lambda}{\sinh \lambda} \notin (-\infty, 0],$$

for all $\lambda \in \mathbb{C}$ with $-\pi < \Im \lambda < \pi$. Let $\nu + ib \in \mathbb{C}$, $|b| < \pi$. Then

$$\begin{aligned} \frac{\nu + ib}{\sinh(\nu + ib)} &= \frac{(\nu + ib) \sinh(\nu - ib)}{|\sinh(\nu + ib)|^2} \\ (4.7) \quad &= \frac{\nu \sinh \nu \cos b + b \cosh \nu \sin b + i(b \sinh \nu \cos b - \nu \cosh \nu \sin b)}{|\sinh(\nu + ib)|^2}. \end{aligned}$$

So

$$\begin{aligned} \frac{\nu + ib}{\sinh(\nu + ib)} \in \mathbb{R} &\Leftrightarrow b \sinh \nu \cos b = \nu \cosh \nu \sin b \\ &\Leftrightarrow b \cot b = \nu \coth \nu \vee \nu = 0. \end{aligned}$$

But $b \cot b \leq 1$ and $\nu \coth \nu > 1$ for $\nu \neq 0$; and $\nu = 0$ means

$$\frac{\nu + ib}{\sinh(\nu + ib)} = \frac{b}{\sin b} > 0.$$

We can conclude

$$(4.8) \quad \frac{\nu + ib}{\sinh(\nu + ib)} \notin (-\infty, 0], \quad \nu \in \mathbb{R}, -\pi < b < \pi.$$

The second inequality is easy, since

$$\begin{aligned} \left| \frac{\nu + ib}{\sinh(\nu + ib)} \right|^2 &= \frac{\nu^2 + b^2}{\sinh^2 \nu + \sin^2 b} \\ &\leq \frac{\nu^2 + b^2}{\sinh^2 \nu} \\ &= \left(1 + \frac{b^2}{\nu^2} \right) \left(\frac{\nu}{\sinh \nu} \right)^2. \end{aligned}$$

To verify the first inequality we show that

$$f(\nu) := \left| \frac{\nu + ib}{\sinh(\nu + ib)} \right|^2 \leq \frac{b^2}{\sin^2 b}, \text{ for all } \nu \in \mathbb{R}.$$

This is obviously true for $\nu = 0$, and since $f(-\nu) = f(\nu)$, we may restrict our analysis to the case $\nu \geq 0$:

$$\begin{aligned} f'(\nu) &= \frac{2\nu(\sinh^2 \nu + \sin^2 b) - (\nu^2 + b^2) \sinh(2\nu)}{(\sinh^2 \nu + \sin^2 b)^2} \\ &= \frac{2\nu \sinh \nu (\sinh \nu - \nu \cosh \nu) + 2\nu \sin^2 b - b^2 \sinh(2\nu)}{(\sinh^2 \nu + \sin^2 b)^2} \\ &\leq 0, \end{aligned}$$

since $\sinh \nu \leq \nu \cosh \nu$ and $2\nu \sin^2 b \leq \sinh(2\nu) \sin^2 b \leq b^2 \sinh(2\nu)$. This means that f stays below $\frac{b^2}{\sin^2 b}$ for all $\nu \in \mathbb{R}$. $\square$

Lemma 4.2. *The function*

$$(4.9a) \quad \psi(\lambda) = \lambda \coth \lambda - a \frac{\lambda}{\sinh \lambda}$$

$$(4.9b) \quad = (1 - a) \lambda \coth \lambda + a \lambda \tanh \frac{\lambda}{2}$$

$$(4.9c) \quad = (1 + a) \lambda \coth \lambda - a \lambda \coth \frac{\lambda}{2}$$

is holomorphic in $\mathbb{C} \setminus \{\pi ki : k \in \mathbb{Z}^*\}$ for $a \neq \pm 1$.

If $a = 1$, then $\psi(\lambda) = \lambda \tanh \frac{\lambda}{2}$ is holomorphic in $\mathbb{C} \setminus \{(2k+1)\pi i : k \in \mathbb{Z}\}$;

if $a = -1$, then $\psi(\lambda) = \lambda \coth \frac{\lambda}{2}$ is holomorphic in $\mathbb{C} \setminus \{2k\pi i : k \in \mathbb{Z}^*\}$.

In particular ψ is holomorphic in the strip $\{z \in \mathbb{C} : |\Im z| < \pi\}$ for any $-1 \leq a \leq 1$.

Proof. The holomorphic properties of ψ follow easily from those of the trigonometric function $\coth$, $\tanh$, and the equalities (4.9), which are easy to verify:

$$\begin{aligned} \psi(\lambda) &= \lambda \coth \lambda - a \frac{\lambda}{\sinh \lambda} \\ &= (1-a)\lambda \coth \lambda + a\lambda \frac{\cosh \lambda - 1}{\sinh \lambda} \\ &= (1-a)\lambda \coth \lambda + a\lambda \tanh \frac{\lambda}{2}, \end{aligned}$$

and

$$\begin{aligned} \psi(\lambda) &= (1+a)\lambda \coth \lambda - a\lambda \frac{\cosh \lambda + 1}{\sinh \lambda} \\ &= (1+a)\lambda \coth \lambda - a\lambda \coth \frac{\lambda}{2}. \end{aligned}$$

□

Lemma 4.3. For $\nu + ib \in \mathbb{C}$, $|b| < \pi$, $\psi(ib)$ is real and

$$(4.10) \quad \Re(\psi(\nu + ib)) \geq \psi(ib),$$

where

$$(4.11a) \quad \psi(ib) = b \cot b - a \frac{b}{\sin b}$$

$$(4.11b) \quad = (1-a)b \cot b + a \left(-b \tan \frac{b}{2} \right)$$

$$(4.11c) \quad = (1+a)b \cot b - ab \cot \frac{b}{2}.$$

Proof. (4.11) is clear. Let $\nu + ib \in \mathbb{C}$, $|b| \leq \pi$, then

$$\begin{aligned} \Re((\nu + ib) \coth(\nu + ib)) &= \frac{1}{2} \frac{\nu \sinh(2\nu) + b \sin(2b)}{\sinh^2 \nu + \sin^2 b} \\ &= \frac{\nu \coth \nu \sinh^2 \nu + b \cot b \sin^2 b}{\sinh^2 \nu + \sin^2 b} \\ &= b \cot b + \sinh^2 \nu \frac{\nu \coth \nu - b \cot b}{\sinh^2 \nu + \sin^2 b} \\ &\geq b \cot b, \end{aligned}$$

and for $|b| \leq \frac{\pi}{2}$:

$$\begin{aligned} \Re((\nu + ib) \tanh(\nu + ib)) &= \frac{1}{2} \frac{\nu \sinh(2\nu) - b \sin(2b)}{\sinh^2 \nu + \cos^2 b} \\ &= \frac{\nu \coth \nu \sinh^2 \nu - b \tan b \cos^2 b}{\sinh^2 \nu + \cos^2 b} \\ &= -b \tan b + \sinh^2 \nu \frac{\nu \coth \nu + b \tan b}{\sinh^2 \nu + \cos^2 b} \\ &\geq -b \tan b. \end{aligned}$$

This gives for $\lambda = \nu + ib \in \mathbb{C}$, $|b| < \pi$:

$$\begin{aligned} \Re(\psi(\lambda)) &= \begin{cases} \Re((1-a)\lambda \coth \lambda + a\lambda \tanh \frac{\lambda}{2}) & , \text{ if } a \geq 0 \\ \Re((1+a)\lambda \coth \lambda - a\lambda \coth \frac{\lambda}{2}) & , \text{ if } a < 0 \end{cases} \\ &\geq \begin{cases} (1-a)b \cot b + a(-b \tan \frac{b}{2}) & , \text{ if } a \geq 0 \\ (1+a)b \cot b - ab \cot \frac{b}{2} & , \text{ if } a < 0 \end{cases} \\ &= \psi(ib). \end{aligned}$$

□

With these preliminary lemmata we can start estimating h by moving the path of integration to $\lambda \mapsto \lambda + ib$:

Lemma 4.4. *There is a constant $C > 0$, such that for all $x, \xi \in \mathbb{R}^n$, $u \in \mathbb{R}$ and $b \in \mathbb{R}$, $|b| < \pi$ the following estimate holds:*

$$(4.12) \quad |h(x, \xi, u)| \leq C \left(\frac{b}{\sin b} \right)^\alpha \exp \left(-2bu - \left(b \cot b - \frac{b}{\sin b} \frac{2x \cdot \xi}{|x|^2 + |\xi|^2} \right) (|x|^2 + |\xi|^2) \right),$$

with $\alpha = \max(0, \frac{n}{2} - 1)$.

Proof. We have

$$\begin{aligned} |h(x, \xi, u)| &= \left| \int_{-\infty}^{\infty} V(\lambda) e^{-\psi(\lambda)R^2 + 2i\lambda u} d\lambda \right| \\ &= \left| \int_{-\infty}^{\infty} V(\lambda + ib) e^{-\psi(\lambda + ib)R^2 + 2i(\lambda + ib)u} d\lambda \right| \\ &\leq \int_{-\infty}^{\infty} |V(\lambda + ib)| e^{-\Re \psi(\lambda + ib)R^2 - 2bu} d\lambda \\ &= e^{-\psi(ib)R^2 - 2bu} \int_{-\infty}^{\infty} |V(\lambda + ib)| d\lambda. \end{aligned}$$

Now we split the remaining integral with $r := \pi - |b|$:

$$\begin{aligned} \int_{-\infty}^{\infty} |V(\lambda + ib)| d\lambda &= 2 \int_0^{\infty} |V(\lambda + ib)| d\lambda \\ &\lesssim \int_0^r \left| \frac{b}{\sin b} \right|^{\frac{n}{2}} d\lambda + \int_r^{\pi} \left(1 + \frac{b^2}{\lambda^2} \right)^{\frac{n}{4}} d\lambda + \int_{\pi}^{\infty} \left| \frac{\lambda}{\sinh \lambda} \right|^{\frac{n}{2}} d\lambda \\ &\lesssim r \left(\frac{b}{\sin b} \right)^{\frac{n}{2}} + \int_r^{\pi} \frac{1}{\lambda^{n/2}} d\lambda + 1 \\ &\lesssim \left(\frac{b}{\sin b} \right)^{\frac{n}{2}-1} + \left(\frac{1}{r} \right)^{\frac{n}{2}-1} + 1 \\ &\lesssim \left(\frac{b}{\sin b} \right)^\alpha, \end{aligned}$$

since

$$(4.13) \quad \frac{b}{\sin b} \sim \frac{1}{\pi - |b|} = \frac{1}{r}.$$

□

Observe that we can assume $u \geq 0$, since $h(x, \xi, -u) = h(x, \xi, u)$.

Setting $b = b_0$, where b_0 parametrizes the corresponding (shortest) geodesic, would not give good results in every case. If b tends to π , the estimate would blow up. To avoid this problem, just set $b = b_0 - \epsilon$ with an appropriate $\epsilon > 0$. (Since we assumed $u \geq 0$, we have $b_0 \geq 0$.)

Proposition 4.1. *For $\zeta = (x, 0)$, $\eta = (\xi, u) \in \mathbb{R}^{n+1}$ we have*

$$(4.14) \quad |h(x, \xi, u)| \lesssim \min \left(1 + \frac{d_{CC}(\zeta, \eta)}{|x + \xi|}, 1 + d_{CC}(\zeta, \eta)^2 \right)^\alpha e^{-d_{CC}(\zeta, \eta)^2},$$

with $\alpha = \max(\frac{n}{2} - 1, 0)$.

Proof. First assume that $x \neq -\xi$. Then the shortest geodesic and therefore the Carnot-Carathéodory distance will be parametrized by $b_0 \in (\pi, \pi)$. We claim that

$$\frac{b_0^2}{\sin^2 b_0} \leq 3 \left(1 + \frac{d_{CC}(\zeta, \eta)^2}{|x + \xi|^2} \right).$$

To see this, first assume that $a < 0$, i.e. $\xi \cdot x < 0$. Then

$$\begin{aligned} d_{CC}(\zeta, \eta)^2 &= \frac{b_0^2}{\sin^2 b_0} (|x|^2 + |\xi|^2 - 2\xi \cdot x \cos b_0) \\ &\geq \frac{b_0^2}{\sin^2 b_0} (|x|^2 + |\xi|^2 + 2\xi \cdot x) \\ &= \frac{b_0^2}{\sin^2 b_0} |x + \xi|^2. \end{aligned}$$

If $a \geq 0$, i.e. $\xi \cdot x \geq 0$, then

$$\begin{aligned} d_{CC}(\zeta, \eta)^2 &\geq \frac{1}{3} \frac{b_0^2}{\sin^2 b_0} (1 - a \cos b_0) (|x|^2 + 2\xi \cdot x + |\xi|^2) \\ &\geq \frac{1}{3} \left(\frac{b_0^2}{\sin^2 b_0} - a \frac{b_0^2 \cos b_0}{\sin^2 b_0} \right) |x + \xi|^2. \end{aligned}$$

If $b_0 > \frac{\pi}{2}$, then $a \frac{b_0^2}{\sin^2 b_0} \cos b_0 < 0$ and if $b_0 \geq \frac{\pi}{2}$, then $a \frac{b_0^2}{\sin^2 b_0} \cos b_0 \leq \frac{b_0^2}{\sin^2 b_0} \leq \frac{\pi^2}{4}$. So

$$\begin{aligned} d_{CC}(\zeta, \eta)^2 &\geq \frac{1}{3} \left(\frac{b_0^2}{\sin^2 b_0} - \frac{\pi^2}{4} \right) |x + \xi|^2 \\ &\geq \frac{1}{3} \left(\frac{b_0^2}{\sin^2 b_0} - 3 \right) |x + \xi|^2. \end{aligned}$$

This proves the claim. Now use Lemma 4.4 with $b = b_0$ to get

$$(4.15) \quad |h(x, \xi, u)| \lesssim \left(1 + \frac{d_{CC}(\zeta, \eta)}{|x + \xi|} \right)^\alpha e^{-d_{CC}(\zeta, \eta)^2}.$$

But this remains true, if $x = -\xi$ (and $x = \xi = 0$), since the right side becomes $+\infty$ in this case. To prove the remaining estimate

$$(4.16) \quad |h(x, \xi, u)| \lesssim (1 + d_{CC}(\zeta, \eta)^2)^\alpha e^{-d_{CC}(\zeta, \eta)^2},$$

we, again, use lemma 4.4. Let $b_0 \in [0, \pi]$ be the parameter for the Carnot-Carathéodory distance. As mentioned above assume that $u \geq 0$. If $b_0 \leq \frac{\pi}{2}$, then

$\frac{b_0}{\sin b_0} \leq \frac{\pi}{2}$, and one can immediately use Lemma 4.4 with $b = b_0$. So assume that $b_0 > \frac{\pi}{2}$. Now set $b = b_0 - \epsilon$ with $\epsilon < \frac{\pi}{2}$, which will be specified later.

If $b_0 \neq \pi$, we have

$$\begin{aligned} b \cot b &= b_0 \cot b_0 + b(\cot b - \cot b_0) - \epsilon \cot b_0 \\ &\geq b_0 \cot b_0 - \epsilon \cot b_0, \end{aligned}$$

since $\cot$ is monotone decreasing on $(0, \pi)$; and

$$\begin{aligned} -b \tan \frac{b}{2} &= -b_0 \tan \frac{b_0}{2} + b \left(\tan \frac{b_0}{2} - \tan \frac{b}{2} \right) + \epsilon \tan \frac{b_0}{2} \\ &\geq -b_0 \tan \frac{b_0}{2} + \epsilon \tan \frac{b_0}{2}, \end{aligned}$$

since $\tan$ is monotone increasing on $(0, \pi)$. Therefore

$$\begin{aligned} \psi(ib) &= \begin{cases} (1-a)b \cot b + a(-b \tan \frac{b}{2}) & , a \geq 0 \\ (1+a)b \cot b - ab \cot \frac{b}{2} & , a < 0 \end{cases} \\ &\geq \psi(ib_0) - \frac{\epsilon}{b_0} \psi(ib_0). \end{aligned}$$

The use of lemma 4.4 with this b yields:

$$\begin{aligned} (4.17) \quad |h(x, \xi, u)| &\lesssim \left(\frac{b}{\sin b} \right)^\alpha e^{-\psi(ib_0)R^2 - 2b_0u + \frac{\epsilon}{b_0}(\psi(ib_0) + 2b_0u)} \\ &\lesssim \left(\frac{1}{\epsilon} \right)^\alpha e^{-(1-\frac{\epsilon}{b_0})d_{CC}(\zeta, \eta)^2}, \end{aligned}$$

first for $b_0 \neq \pi$. But, if $b_0 = \pi$, then we have $x = \xi = 0$ or $x = -\xi$. We claim that in this case estimate (4.17) remains true: If $x = \xi = 0$ then

$$\begin{aligned} |h(x, \xi, u)| &\lesssim \left(\frac{b}{\sin b} \right)^\alpha e^{-2b_0u + 2b_0u \frac{\epsilon}{b_0}} \\ &\lesssim \left(\frac{1}{\epsilon} \right)^\alpha e^{-(1-\frac{\epsilon}{b_0})d_{CC}(\zeta, \eta)^2}, \end{aligned}$$

since $2b_0u = 2\pi u = d_{CC}(\zeta, \eta)^2$. And, if $x = -\xi$, then $a = -1$ and $\psi(ib) = b \cot \frac{b}{2} \geq 0$, so that

$$\begin{aligned} |h(x, \xi, u)| &\lesssim \left(\frac{b}{\sin b} \right)^\alpha e^{-\psi(ib)R^2 - 2b_0u + 2b_0u \frac{\epsilon}{b_0}} \\ &\lesssim \left(\frac{1}{\epsilon} \right)^\alpha e^{-2b_0u + 2b_0u \frac{\epsilon}{b_0}} \\ &= \left(\frac{1}{\epsilon} \right)^\alpha e^{-(1-\frac{\epsilon}{b_0})d_{CC}(\zeta, \eta)^2}, \end{aligned}$$

since again $2b_0u = 2\pi u = d_{CC}(\zeta, \eta)^2$.

Setting $\epsilon = \frac{1}{1+d_{CC}(\zeta, \eta)^2}$, which optimizes this inequality (up to a constant), gives us the desired result:

$$(4.18) \quad |h(x, \xi, u)| \lesssim (1 + d_{CC}(\zeta, \eta)^2)^\alpha e^{-d_{CC}(\zeta, \eta)^2}.$$

□

This gives the following main result:

Theorem 4.1. *For $\zeta = (x, 0)$, $\eta = (\xi, u) \in \mathbb{R}^{n+1}$ we have*

$$(4.19) \quad |K_t(x, \xi, u)| \lesssim t^{-\frac{n}{2}-1} \min \left(1 + \frac{d_{CC}(\zeta, \eta)}{|x + \xi|}, 1 + \frac{d_{CC}(\zeta, \eta)^2}{4t} \right)^\alpha e^{-\frac{1}{4t}d_{CC}(\zeta, \eta)^2},$$

with $\alpha = \max\left(\frac{n}{2} - 1, 0\right)$.

Proof. The proof is almost trivial, if one uses (4.2) and the homogeneity of d_{CC} . $\square$

REFERENCES

- [1] Richard Beals, Bernard Gaveau, and Peter C. Greiner. Hamilton-Jacobi theory and the heat kernel on Heisenberg groups. *J. Math. Pures Appl. (9)*, 79(7):633–689, 2000.
 - [2] Ovidiu Calin and Der-Chen Chang. The geometry on a step 3 Grushin model. *Appl. Anal.*, 84(2):111–129, 2005.
 - [3] Ovidiu Calin, Der-Chen Chang, Peter Greiner, and Yakar Kannai. On the geometry induced by a Grusin operator. In *Complex analysis and dynamical systems II*, volume 382 of *Contemp. Math.*, pages 89–111. Amer. Math. Soc., Providence, RI, 2005.
 - [4] W.L. Chow. Über Systeme von Linearen Partiellen Differentialgleichungen erster Ordnung. *Math. Ann.*, 117:98–105, 1939.
 - [5] Gerald B. Folland. *Harmonic analysis in phase space*, volume 122 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1989.
 - [6] Peter C. Greiner, David Holcman, and Yakar Kannai. Wave kernels related to second-order operators. *Duke Math. J.*, 114(2):329–386, 2002.
 - [7] Ralf Meyer. *L^p -estimates for the wave equation associated to the Grusin operator*. PhD thesis, Kiel, Univ., 2006.
 - [8] Adam Sikora. Riesz transform, Gaussian bounds and the method of wave equation. *Math. Z.*, 247(3):643–662, 2004.
 - [9] Elias M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1993. With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III.
 - [10] Robert S. Strichartz. Sub-Riemannian geometry. *J. Differential Geom.*, 24(2):221–263, 1986.
 - [11] Robert S. Strichartz. Corrections to: “Sub-Riemannian geometry” [*J. Differential Geom.* **24** (1986), no. 2, 221–263; MR0862049 (88b:53055)]. *J. Differential Geom.*, 30(2):595–596, 1989.
- E-mail address:* martin@inter-zone.de